arXiv:1603.04076v1 [math.NT] 13 Mar 2016

TWISTED CHARACTERISTIC p ZETA FUNCTIONS

BRUNO ANGLÈS, TUAN NGO DAC, AND FLORIC TAVARES RIBEIRO

Abstract. We propose a "twisted" variation of zeta functions introduced by David Goss in 1979.

Contents

1. Introduction 1
2. Notation and preliminaries 3
3. A Key Lemma 7
4. Several variable twisted zeta functions 12
4.1. The ∞-adic case 13
4.2. The case of finite places 16
5. Examples 17
5.1. The case $A = \mathbb{F}_q[\theta]$ 18
5.2. Twisted Goss zeta functions 21
5.3. A-harmonic series attached to some admissible maps 22
6. Multiple several variable twisted zeta functions 24
References 28

1. Introduction

Let $\mathbb{F}_q$ be a finite field having q elements, and let θ be an indeterminate over $\mathbb{F}_q$; the Carlitz-Goss zeta function is defined as follows ([21], chapter 8):

$$\zeta_A((x;y)) := \sum_{d\geq 0}\Big(\sum_{a\in A_{+,d}} \frac{1}{(\frac{a}{\theta^d})^y}\Big)x^{-d} \in \mathbb{C}_\infty^\times,$$

where $(x;y) \in \mathbb{C}_\infty^\times \times \mathbb{Z}_p$, $A := \mathbb{F}_q[\theta]$, $A_{+,d} := \{a \in A, a \,\text{monic}\,, \deg_\theta a = d\}$, and $\mathbb{C}_\infty$ is the completion of an algebraic closure of $\mathbb{F}_q((\frac{1}{\theta}))$. The facts that $\zeta_A(.)$ converges on $\mathbb{S}_\infty := \mathbb{C}_\infty^\times \times \mathbb{Z}_p$ and is "essentially algebraic" (i.e. for $y \in \mathbb{N}$, $\zeta_A((\theta^y x; -y))$ is a polynomial in x^{-1} with coefficients in A) can be proved by using the following vanishing result: for $n \in \mathbb{N}$, for $d(q-1) > \ell_q(n)$, $\sum_{a\in A_{+,d}} a^n = 0$, where $\ell_q(n)$ is the sum of digits of n in base q (this is a consequence of the combinatorial result: [21], Lemma 8.8.1). The Carlitz-Goss zeta function is an example of a new kind of L-series introduced by D. Goss in [19] (see also [20]). The "special values" of these type of L-functions are at the heart of function field arithmetic and we refer the interested reader to (this list is clearly not exhaustive): [3], [13], [24], [36].

Date: March 15, 2016.

Let $s, n \in \mathbb{N}$, $n \geq 1$, let $x \in \mathbb{C}_\infty^\times$ and $y_1, \dots, y_n \in \mathbb{Z}_p$; let's consider the following "twisted" version of the Carlitz-Goss zeta function:

$$\zeta_A(t_1, \dots, t_s; \theta_1, \dots, \theta_n; (x; y_1, \dots, y_n)) := \sum_{d \geq 0} \Big(\sum_{a \in A_{+,d}} \frac{a(t_1) \cdots a(t_s)}{(\frac{a(\theta_1)}{\theta_1^d})^{y_1} \cdots (\frac{a(\theta_n)}{\theta_n^d})^{y_n}} \Big) x^{-d},$$

where $t_1, \dots, t_s \in \mathbb{C}_\infty$, $\theta_1, \dots, \theta_n \in \mathbb{C}_\infty^\times$ and are such that $\frac{1}{\theta_i}, i = 1, \dots, n$, is in the maximal ideal of the valuation ring of $\mathbb{C}_\infty$. Using the fact that $\forall t_1, \dots, t_s \in \mathbb{C}_\infty$, $\forall d > \frac{s}{q-1}$, $\sum_{a \in A_{+,d}} a(t_1) \cdots a(t_s) = 0$ (this is again a consequence of [21], Lemma 8.8.1), one can prove that $\zeta_A(t_1, \dots, t_s; \theta_1, \dots, \theta_n; .)$ converges on $\mathbb{C}_\infty^\times \times \mathbb{Z}_p^n$ and that this function is essentially algebraic, i.e. for $y_1, \dots, y_n \in \mathbb{N}$:

$$\zeta_A(t_1, \dots, t_s; \theta_1, \dots, \theta_n; (\prod_{i=1}^n \theta_i^{y_i} x; -y_1, \dots, -y_n)) \in \mathbb{F}_q[t_1, \dots, t_s, \theta_1, \dots, \theta_n, x^{-1}].$$

Observe that if $s = 0, n = 1$ and $\theta_1 = \theta$, we recover the Carlitz-Goss zeta function, and if $s \geq 0$, $n = 1$, $\theta_1 = \theta$, we recover the L-series introduced by F. Pellarin in [28].

Our aim in this article is to extend the above construction to the case where $K/\mathbb{F}_q$ is a global function field, ∞ is a place of K, and A is the ring of elements of K which are regular outside ∞. Although the twisted zeta functions proposed in this paper are in the spirit of the twisted Carlitz-Goss zeta functions defined above, the proof of the convergence of these functions, and their v-adic interpolation at finite places v of K, are more subtle and based on a technical key Lemma which generalizes the vanishing result mentioned above (Lemma 3.2). The zeta functions introduced by D. Goss as well as their deformations over affinoid algebras are examples of such twisted zeta functions (see example 5.2).

The main ingredient to our construction is what we called *admissible maps* (section 2). Let K_∞ be the ∞-adic completion of K, let $\mathbb{F}_\infty$ be the residue field of K_∞, and let $\mathrm{sgn} : K_\infty^\times \to \mathbb{F}_\infty^\times$ be a "sign function", i.e. sgn is a group homomorphism such that $\mathrm{sgn}\,|_{\mathbb{F}_\infty^\times} = \mathrm{Id}_{\mathbb{F}_\infty^\times}$. Let π be a prime of K_∞ such that $\pi \in \mathrm{Ker}\,\mathrm{sgn}$. Let $\overline{K}_\infty$ be a fixed algebraic closure of K_∞, let $\overline{\mathbb{F}}_q$ be the algebraic closure of $\mathbb{F}_q$ in $\overline{K}_\infty$, then sgn can be naturally extended to a morphism $\mathrm{sgn} : \overline{K}_\infty \to \overline{\mathbb{F}}_q^\times$ (see section 2) according to our choice of π. Let $v_\infty : \overline{K}_\infty \to \mathbb{Q} \cup \{+\infty\}$ be the valuation on $\overline{K}_\infty$ such that $v_\infty(\pi) = 1$. For $x \in \overline{K}_\infty^\times$, let's set:

$$\langle x \rangle = \frac{x}{\mathrm{sgn}(x) \pi^{v_\infty(x)}}.$$

Let $\mathcal{I}(A)$ be the group of non-zero fractional ideals of A. By definition, an admissible map $\eta : \mathcal{I}(A) \to \overline{K}_\infty$ is a map such that there exist $n(\eta) \in \mathbb{Z}_p$, $\gamma(\eta) \in \overline{K}_\infty^\times$, and an open subgroup of finite index $M(\eta) \subset K_\infty^\times$, with the following property:

$$\forall I \in \mathcal{I}(A), \forall a \in K^\times \cap M(\eta), \quad \eta(Ia) = \eta(I) \langle a \rangle^{n(\eta)} \gamma_\eta^{v_\infty(a)}.$$

The twisted zeta functions considered in this work are constructed with the help of such admissible maps (section 4). Let's give a basic example. Let $\eta : \mathcal{I}(A) \to \overline{K}_\infty^\times$ be an admissible map such that $n(\eta) = 1$ and η is algebraic (i.e. $\eta(\mathcal{I}(A)) \subset \overline{K}^\times$, where $\overline{K}$ is the algebraic closure of K in $\overline{K}_\infty$). Let's set:

$$K(\eta) = K(\mathbb{F}_\infty, \gamma_\eta \pi^{-1}, \eta_i(I), I \in \mathcal{I}(A)),$$
$$K_\infty(\langle \eta \rangle) = K_\infty(\langle \gamma_\eta \rangle, \langle \eta_i(I) \rangle, I \in \mathcal{I}(A)).$$

Observe that $K(\eta)/K$ is a finite extension, and $K_\infty(\langle\eta\rangle)/K_\infty$ is also a finite extension. Let $s, n \in \mathbb{N}$, $n \geq 1$. Let $\mathbb{C}_\infty$ be the completion of $\overline{K}_\infty$. Let $\rho_1, \dots \rho_s : K(\eta) \hookrightarrow \mathbb{C}_\infty$ be s $\mathbb{F}_p$-algebra homomorphisms, and let $\sigma_1, \dots, \sigma_n : K_\infty(\langle\eta\rangle) \hookrightarrow \mathbb{C}_\infty$ be n continuous $\mathbb{F}_p$-algebra homomorphisms. For $u = (x; (y_1, \dots, y_n)) \in \mathbb{C}_\infty^\times \times \mathbb{Z}_p^n$, let's set:

$$\zeta_{\eta,A}(\underline{\rho}; \underline{\sigma}; u) = \sum_{d \geq 0} \Big(\sum_{\substack{I \in \mathcal{I}(A), I \subset A \\ \deg I = d}} \frac{\prod_{i=1}^s \rho_i(\eta(I))}{\prod_{j=1}^n \sigma_j(\langle\eta(I)\rangle^{-y_j})} \Big) x^{-d}.$$

Then, as a consequence of Theorem 4.1, $\zeta_{\eta,A}(\underline{\rho}; \underline{\sigma}; .)$ converges on $\mathbb{C}_\infty^\times \times \mathbb{Z}_p^n$. One can easily notice that the above function, and its twists by some characters, generalize the twisted Carlitz-Goss zeta function. We refer the reader to section 5 for more detailed examples.

For the sake of completeness, in the last section of this article, we also introduce multiple twisted zeta functions, as well as their v-adic interpolation at finite places v of K, in the spirit of Thakur's multiple zeta values ([4],[39]).

2. Notation and preliminaries

Let $\mathbb{F}_q$ be a finite field having q elements and of characteristic p. Let $K/\mathbb{F}_q$ be a global function field ($\mathbb{F}_q$ is algebraically closed in K). Let ∞ be a place of K. We denote by K_∞ the ∞-adic completion of K, and by $\mathbb{F}_\infty$ the residue field of K_∞. We fix $\overline{K}_\infty$ an algebraic closure of K_∞. Let $\overline{K}$ be the algebraic closure of K in $\overline{K}_\infty$, and let $\overline{\mathbb{F}}_q \subset \overline{K}$ be the algebraic closure of $\mathbb{F}_q$ in $\overline{K}_\infty$. Let $\mathbb{C}_\infty$ be the completion of $\overline{K}_\infty$. Let A be the ring of elements of K which are regular outside of ∞. Let $v_\infty : K_\infty \to \mathbb{Z} \cup \{+\infty\}$ be the normalized discrete valuation associated to the local field K_∞. We still denote by $v_\infty : \mathbb{C}_\infty \to \mathbb{R} \cup \{+\infty\}$ the extension of v_∞ to $\mathbb{C}_\infty$.

Let $\mathcal{I}(A)$ be the group of fractional ideals of A, and $\mathcal{P}(A) = \{\alpha A, \alpha \in K^\times\}$. Recall that $\mathrm{Pic}(A) = \frac{\mathcal{I}(A)}{\mathcal{P}(A)}$ is a finite abelian group. For any ideal $I \subset A, I \neq \{0\}$, we set:

$$\deg I = \dim_{\mathbb{F}_q} \frac{A}{I}.$$

Note that this function on non-zero ideals of A extends naturally to a morphism $\deg : \mathcal{I}(A) \to \mathbb{Z}$. In particular, we have:

$$\forall \alpha \in K^\times, \quad \deg \alpha := \deg \alpha A = -d_\infty v_\infty(\alpha),$$

where $d_\infty = \dim_{\mathbb{F}_q} \mathbb{F}_\infty$.

Let $x \in \overline{K}_\infty$ such that $v_\infty(x) > 0$. Let $(x_n)_{n \geq 1}$ be a sequence of elements in $\overline{K}_\infty$ such that $x_1 = x$, and for $n \geq 1$, $x_{n+1}^{n+1} = x_n$. Let $z \in \mathbb{Q}$, write $z = \frac{m}{n!}, m \in \mathbb{Z}, n \geq 1$, we set:

$$x^z = x_n^m.$$

This is well-defined. We set:

$$U_\infty = \{y \in \overline{K}_\infty, v_\infty(y-1) > 0\}.$$

Observe that U_∞ is a $\mathbb{Q}_p$-vector space. We have:

$$\overline{K}_\infty^\times = x^{\mathbb{Q}} \times \overline{\mathbb{F}}_q^\times \times U_\infty.$$

Let $y \in \overline{K}_\infty^\times$, then y can be written in a unique way:

$$y = x^{\frac{v_\infty(y)}{v_\infty(x)}} \omega_x(y) \langle y \rangle_x, \quad \omega_x(y) \in \overline{\mathbb{F}}_q^\times, \langle y \rangle_x \in U_\infty.$$

Observe that the map $\langle . \rangle_x : \overline{K}_\infty^\times \to U_\infty$ is a group homomorphism such that for $y \in U_\infty$, $\langle y \rangle_x = y$. We will need the following Lemma in the sequel (see also [22], Lemma 2):

Lemma 2.1.
1) Let $\langle . \rangle' : \overline{K}_\infty^\times \to U_\infty$ be a group homomorphism such that for $\forall y \in U_\infty, \langle y \rangle' = y$. Then there exists a unique element $u \in U_\infty$ such that:

$$\forall y \in \overline{K}_\infty^\times, \quad \langle y \rangle' = \langle y \rangle_x u^{v_\infty(y)}.$$

2) Let $\mathrm{sgn} : K_\infty^\times \to \overline{\mathbb{F}}_q^\times$ be a group homomorphism such that $\forall \zeta \in \mathbb{F}_q^\times, \mathrm{sgn}(\zeta) = \zeta$. Then there exist $e \in \mathbb{N}$, $e \equiv 1 \pmod{q-1}$, and an element $\zeta' \in \overline{\mathbb{F}}_q^\times$, such that:

$$\forall y \in K_\infty^\times, \quad \mathrm{sgn}(y) = \omega_x(y)^e \zeta'^{v_\infty(y)}.$$

3) For $i = 1, 2$, let $\langle . \rangle_i : \overline{K}_\infty^\times \to U_\infty$ be a group homomorphism such that $\forall y \in U_\infty, \langle y \rangle_i = y$, and let $\gamma_i \in \overline{K}_\infty^\times$. Let's assume that there exist an open subgroup of finite index $N \subset K_\infty^\times$, and $n_1, n_2 \in \mathbb{Z}_p$, such that:

$$\forall a \in N \cap K^\times, \quad \langle a \rangle_1^{n_1} \gamma_1^{v_\infty(a)} = \langle a \rangle_2^{n_2} \gamma_2^{v_\infty(a)}.$$

Then $n_1 = n_2$.

Proof.
1) We have:

$$u = (\langle x \rangle')^{\frac{1}{v_\infty(x)}}.$$

2) There exists $e \in \mathbb{N}$, $e \equiv 1 \pmod{q-1}$ such that:

$$\forall \zeta \in \mathbb{F}_\infty^\times, \quad \mathrm{sgn}(\zeta) = \zeta^e.$$

Let $U = K_\infty^\times \cap U_\infty$. We get:

$$\frac{\mathrm{sgn}(.)}{\omega_x(.)^e} \mid_{\mathbb{F}_\infty^\times \times U} = 1.$$

Let π be any element of $K_\infty^\times$ such that $v_\infty(\pi) = 1$, and set:

$$\zeta' = \frac{\mathrm{sgn}(\pi)}{\omega_x(\pi)^e} \in \overline{\mathbb{F}}_q^\times.$$

3) Let $a \in U_\infty \cap N \cap K^\times$, $a \neq 1$. Then:

$$a^{n_1} = a^{n_2}.$$

Thus $n_1 = n_2$.

□

A map $\eta : \mathcal{I}(A) \to \overline{K}_\infty$ will be called *admissible* if there exist an open subgroup of finite index $N \subset K_\infty^\times$, an element $n(\eta) \in \mathbb{Z}_p$, an element $\gamma \in \overline{K}_\infty^\times$, and a group homomorphism $\langle . \rangle : \overline{K}_\infty \to U_\infty$, with $\langle . \rangle \mid_{U_\infty} = \mathrm{Id}_{U_\infty}$, such that:

$$\forall I \in \mathcal{I}(A), \forall \alpha \in K^\times \cap N, \quad \eta(I\alpha) = \eta(I) \langle \alpha \rangle^{n(\eta)} \gamma^{v_\infty(\alpha)}.$$

By Lemma 2.1, this definition does not depend on the choice of $\langle . \rangle$, $n(\eta)$ is well-defined, and γ is well-defined modulo $U_\infty^{n(\eta)}$. Note also that the product of admissible maps is an admissible map.

We observe that any constant map from $\mathcal{I}(A)$ to $\overline{K}_\infty$ is admissible. An admissible map η will be called algebraic if $\eta(\mathcal{I}(A)) \subset \overline{K}$.

Let's give a fundamental example, due to D. Goss, of such an admissible map ([21], section 8.2). Let t be a prime of K_∞ (i.e. $v_\infty(t) = 1$). For $x \in K_\infty^\times$, let's write:

$$x = t^{v_\infty(x)} \omega_t(x) \langle x \rangle_t, \quad \omega_t(x) \in \mathbb{F}_\infty^\times, v_\infty(\langle x \rangle_t - 1) > 0.$$

Write $d_\infty = p^k \bar{d}_\infty$ with $\bar{d}_\infty \not\equiv 0 \pmod p$. Let p^m be the biggest power of p that divides $\mid \mathrm{Pic}(A) \mid$. Set $l = \mathrm{Max}\{m, k\}$. We fix $t^{\frac{1}{d_\infty}} \in \overline{K}_\infty$ a $\bar{d}_\infty$-th root of t. Let:

$$T = \mathbb{F}_\infty((t^{\frac{1}{p^l d_\infty}})).$$

If $I \in \mathcal{I}(A)$, then there exists an integer h dividing $\mid \mathrm{Pic}(A) \mid$, such that:

$$I^h = \alpha A, \quad \alpha \in K^\times.$$

We set:

$$[I]_t = (\langle \alpha \rangle_t)^{\frac{1}{h}} t^{\frac{-deg I}{d_\infty}} \in T.$$

Then, $K([I], I \in \mathcal{I}(A))/K$ is a finite extension, $[.]_t : \mathcal{I}(A) \to \overline{K}_\infty^\times$ is a group homomorphism and:

$$\forall \alpha \in K^\times, \quad [\alpha A]_t = \frac{\alpha}{\omega_t(\alpha)},$$

Thus, by Lemma 2.1, $[.]_t$ is an algebraic admissible map.

From now in, we fix a prime π of K_∞ ($\pi \in K_\infty$, $v_\infty(\pi) = 1$). We also fix a sequence $(\pi_n)_{n \geq 1}$ of elements in $\overline{K}_\infty$ such that $\pi_1 = \pi$, and $\forall n \geq 1$, $\pi_{n+1}^{n+1} = \pi_n$. By our previous construction, attached to this choice of such a sequence, we have a group homomorphism $\langle . \rangle = \langle . \rangle_\pi : \overline{K}_\infty^\times \to U_\infty$, and a group homomorphism $\mathrm{sgn} = \omega_\pi : \overline{K}_\infty^\times \to \overline{\mathbb{F}}_q^\times$. For $I \in \mathcal{I}(A)$, we set:

$$[I] = \langle I \rangle \pi^{-\frac{\deg I}{d_\infty}} \in \pi^{\mathbb{Q}} \times U_\infty,$$

where:

$$\langle I \rangle = \langle \alpha \rangle^{\frac{1}{h}}, \quad I^h = \alpha A, \quad \alpha \in K^\times.$$

Then, $[.] : \mathcal{I}(A) \to \overline{K}_\infty^\times$ is a group homomorphism such that:

$$\forall \alpha \in K^\times, \quad [\alpha A] = \frac{\alpha}{\mathrm{sgn}(\alpha)}.$$

Observe that $[.]$ is an algebraic admissible map, and $\langle . \rangle : \mathcal{I}(A) \to U_\infty$ is an admissible map.

Let $\eta : \mathcal{I}(A) \to \overline{K}_\infty$ be an admissible map. Then there exist $M(\eta) \subset K_\infty^\times$ an open subgroup of finite index, an element $n(\eta) \in \mathbb{Z}_p$, an element $\gamma_\eta \in \overline{K}_\infty^\times$, such that:

$$\forall I \in \mathcal{I}(A), \forall \alpha \in K^\times \cap M(\eta), \quad \eta(I\alpha) = \eta(I)(\frac{\alpha}{\mathrm{sgn}(\alpha)\pi^{v_\infty(\alpha)}})^{n(\eta)} \gamma_\eta^{v_\infty(\alpha)}.$$

Let's observe that, if $n(\eta) \in \mathbb{Z}$, $\forall \alpha \in K^\times \cap M(\eta)$, we have:

$$(\frac{\alpha}{\mathrm{sgn}(\alpha)\pi^{v_\infty(\alpha)}})^{n(\eta)}\gamma_\eta^{v_\infty(\alpha)} = (\frac{\alpha}{\mathrm{sgn}(\alpha)})^{n(\eta)}(\gamma_\eta\pi^{-n(\eta)})^{v_\infty(\alpha)}.$$

In that case, we set $K(\eta) := K(\mathbb{F}_\infty, \gamma_\eta\pi^{-n(\eta)}, \eta(I), I \in \mathcal{I}(A))$. Let's observe that, if η is algebraic then $K(\eta)/K$ is a finite extension.

Let (F, v_F) be a field which is complete with respect to a non-trivial valuation $v_F : F \to \mathbb{R} \cup \{+\infty\}$ and such that F is a $\overline{\mathbb{F}}_q$-algebra.

Let $n \geq 1$, we set:

$$\mathbb{S}_{F,n} = F^\times \times \mathbb{Z}_p^n.$$

We view $\mathbb{S}_{F,n}$ as an abelian group with group action written additively. Let $\eta_1, \dots, \eta_n : \mathcal{I}(A) \to \overline{K}_\infty^\times$ be n admissible maps (note that for $i = 1, \dots, n$, $\forall I \in \mathcal{I}(A)$, $\eta_i(I) \neq 0$). Observe that for $i = 1, \dots, n$, $K_\infty(\langle\eta_i\rangle) := K_\infty(\langle\gamma_{\eta_i}\rangle, \langle\eta_i(I)\rangle, I \in \mathcal{I}(A))$ is a finite extension of K_∞. For $i = 1, \dots, n$, let $\sigma_i : K_\infty(\langle\eta_i\rangle) \hookrightarrow F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. For $u = (x; z_1, \dots, z_n) \in \mathbb{S}_{F,n}$ and $I \in \mathcal{I}(A)$, we set:

$$I^{\underline{z}}_{\underline{\sigma},\underline{\eta}} = \prod_{i=1}^n \sigma_i(\langle\eta_i(I)\rangle^{z_i}) \in F^\times,$$

$$I^u_{\underline{\sigma},\underline{\eta}} = x^{\deg I}\prod_{i=1}^n \sigma_i(\langle\eta_i(I)\rangle^{z_i}) \in F^\times.$$

Observe that for $I \in \mathcal{I}(A)$, $u, u' \in \mathbb{S}_{F,n}$ and $\alpha \in K^\times \cap M(\eta_1) \cap \cdots \cap M(\eta_n)$, we have:

$$I^{u+u'}_{\underline{\sigma},\underline{\eta}} = I^u_{\underline{\sigma},\underline{\eta}} I^{u'}_{\underline{\sigma},\underline{\eta}},$$

$$(I\alpha)^u_{\underline{\sigma},\underline{\eta}} = I^u_{\underline{\sigma},\underline{\eta}}(\alpha A)^u_{\underline{\sigma},\underline{\eta}},$$

where:

$$(\alpha A)^u_{\underline{\sigma},\underline{\eta}} := (x^{-d_\infty}\prod_{i=1}^n \sigma_i(\langle\gamma_{\eta_i}\rangle^{z_i}))^{v_\infty(\alpha)}\prod_{i=1}^n \sigma_i(\langle\alpha\rangle^{n(\eta_i)z_i}) \in F^\times.$$

Remark 2.2. *Let $N \subset K_\infty^\times$ be an open subgroup of finite index, and let's set:*

$$\mathcal{P}(N) = \{xA, x \in K^\times \cap N\}.$$

Then $\frac{\mathcal{I}(A)}{\mathcal{P}(N)}$ is a finite abelian group. Let $\chi : \mathcal{I}(A) \to \overline{K}_\infty$ be a map such that:

$$\forall I \in \mathcal{I}(A), \forall J \in \mathcal{P}(N), \quad \chi(IJ) = \chi(I).$$

Observe that χ is an admissible map. Let $\gamma' \in \overline{K}_\infty^\times$, and let $z \in \mathbb{Z}_p$. Then

$$\eta = \chi(.)\langle.\rangle^z(\gamma')^{\deg(.)}$$

is an admissible map with $M(\eta) = N$, $n(\eta) = z$, and $\gamma_\eta = (\gamma')^{-d_\infty}$.

Reciprocally, let η be an admissible map. Select $\gamma' \in \overline{K}_\infty^\times$ such that $(\gamma')^{d_\infty} = \frac{1}{\gamma_\eta}$. Set $\chi(.) = \frac{\eta(.)}{\langle.\rangle^{n(\eta)}(\gamma')^{\deg(.)}}$. Then $\chi : \mathcal{I}(A) \to \overline{K}_\infty$ is such that:

$$\forall I \in \mathcal{I}(A), \forall J \in \mathcal{P}(M(\eta)), \quad \chi(IJ) = \chi(I).$$

Remark 2.3. *Let T/K_∞ be a finite extension, let $\mathbb{F}_T$ be the residue field of T, and let π' be a prime of T. Let $\sigma : T \hookrightarrow F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. Let $\varphi \in \mathrm{Gal}(T/\mathbb{F}_p((\pi'))) \simeq \mathrm{Gal}(\mathbb{F}_T/\mathbb{F}_p)$, such that:*

$$\forall \zeta \in \mathbb{F}_T, \quad \sigma(\zeta) = \varphi(\zeta).$$

Let $y = \sigma(\pi')$. Then $v_F(y) > 0$, and:

$$\forall a \in T, \quad \sigma(a) = \varphi(a) \mid_{\pi'=y} .$$

Thus the choice of a continuous $\mathbb{F}_p$-algebra homomorphism $\sigma : T \hookrightarrow F$ is equivalent to the choice of an element $y \in F^\times$, $v_F(y) > 0$, and $\varphi \in \mathrm{Gal}(\mathbb{F}_T/\mathbb{F}_p)$.

3. A Key Lemma

Let E/K be a finite extension, $E \subset \overline{K}$, let g_E be the genus of E, and let O_E be the integral closure of A in E. Let $\mathcal{I}(O_E)$ be the group of fractional ideals of O_E, and let $\mathcal{P}(O_E) = \{xO_E, x \in E^\times\}$. Let $N_{E/K} : \mathcal{I}(O_E) \to \mathcal{I}(A)$ be the group homomorphism such that, if $\mathfrak{P}$ is a maximal ideal of O_E and if $P = \mathfrak{P} \cap A$ is the corresponding maximal ideal of A, then:

$$N_{E/K}(\mathfrak{P}) = P^{[\frac{O_E}{\mathfrak{P}} : \frac{A}{P}]}.$$

Note that, if $\mathfrak{I} = xO_E, x \in E^\times$, then:

$$N_{E/K}(\mathfrak{I}) = N_{E/K}(x)A,$$

where $N_{E/K} : E \to K$ denotes also the usual norm map. Let $\infty_1, \dots, \infty_r$ be the places of E above ∞. For $i = 1, \dots, r$, let f_i be the residual degree of ∞_i/∞. Let $v_1, \dots, v_r$ be the valuations on E associated to these places. We denote by E_{v_i} the v_i-adic completion of E, $i = 1, \dots, r$.

Let $\mathcal{B} \subset O_E$, $\mathcal{B} \in \mathcal{I}(O_E)$. We denote by $\mathcal{I}(\mathcal{B})$ the group of fractional ideals of O_E which are relatively prime to $\mathcal{B}$. For $i = 1, \dots, r$, let N_i be an open subgroup of finite index of $E_{v_i}^\times$, and let $m_i \geq 0$ be the least integer such that:

$$\{x \in E_{v_i}^\times, v_i(x-1) \geq m_i\} \subset N_i.$$

We set $N = \prod_{i=1}^r N_i$, and:

$$\mathcal{P}(\mathcal{B}, N) = \{xO_E \in \mathcal{I}(\mathcal{B}), x \in E^\times, x \equiv 1 \pmod{\mathcal{B}}, x \in N_i, i = 1, \dots, r\}.$$

Observe that $\frac{\mathcal{I}(\mathcal{B})}{\mathcal{P}(\mathcal{B},N)}$ is a finite abelian group. Let:

$$d(\mathcal{B}) = \dim_{\mathbb{F}_q} \frac{O_E}{\mathcal{B}},$$

$$d_\infty(N) = \sum_{i=1}^r d_\infty f_i(m_i + 1).$$

More generally, let $\mathcal{S}$ be a finite set, possibly empty, of maximal ideals of O_E which are relatively prime to $\mathcal{B}$. We denote by $\mathcal{I}_\mathcal{S}(\mathcal{B})$ the group of fractional ideals of $\mathcal{I}(O_E)$ which are relatively prime to $\mathcal{B}\prod_{\mathfrak{P}\in\mathcal{S}}\mathfrak{P}$. We also set $\mathcal{P}_\mathcal{S}(\mathcal{B}, N) = \mathcal{P}(\mathcal{B}, N) \cap \mathcal{I}_\mathcal{S}(\mathcal{B})$. Observe that we have a natural group isomorphism:

$$\frac{\mathcal{I}_\mathcal{S}(\mathcal{B})}{\mathcal{P}_\mathcal{S}(\mathcal{B}, N)} \simeq \frac{\mathcal{I}(\mathcal{B})}{\mathcal{P}(\mathcal{B}, N)}.$$

We set:

$$d_{\mathcal{S}}(\mathcal{B}) = \dim_{\mathbb{F}_q} \frac{O_E}{\mathcal{B}\prod_{\mathfrak{P}\in\mathcal{S}}\mathfrak{P}}.$$

We recall a basic result ([21], Lemma 8.8.1):

Lemma 3.1. *Let $r \in \mathbb{N}$, and let W be a finite dimensional $\mathbb{F}_p$-vector space. For $i = 1, \dots, r$, let $f_i : W \to F$ be an $\mathbb{F}_p$-linear map. If $\dim_{\mathbb{F}_p} W > \frac{r}{p-1}$, then:*

$$\forall x_1, \dots, x_r \in F, \qquad \sum_{w\in W} (x_1 + f_1(w)) \cdots (x_r + f_r(w)) = 0.$$

Proof. We recall its proof for the convenience of the reader. First, note that for $k, m_1, \dots m_k \in \mathbb{N}$, $k \geq 1$, then :

$$\sum_{\zeta_1,\dots,\zeta_k\in\mathbb{F}_p} \zeta_1^{m_1} \cdots \zeta_k^{m_k} \neq 0 \Leftrightarrow \forall i \in \{1, \dots k\}, \quad m_i \equiv 0 \pmod{p-1} \text{and} \, m_i \geq 1.$$

Thus, in particular, if $\sum_{\zeta_1,\dots,\zeta_k\in\mathbb{F}_p} \zeta_1^{m_1} \cdots \zeta_k^{m_k} \neq 0$ then $m_1 + \cdots + m_k \geq k(p-1)$. Let $\alpha = \dim_{\mathbb{F}_p} W$. Let's assume that $\alpha > \frac{r}{p-1}$. Let

$$S = \sum_{w\in W} (x_1 + f_1(w)) \cdots (x_r + f_r(w)).$$

Let's set:

$$W = \oplus_{j=1}^{\alpha} \mathbb{F}_p e_j.$$

Then:

$$S = \sum_{\zeta_1,\dots,\zeta_\alpha\in\mathbb{F}_p} \prod_{j=1}^{r} (x_j + \zeta_1 f_j(e_1) + \cdots + \zeta_\alpha f_j(e_\alpha)).$$

When we develop the above product and sum over $\zeta_1, \dots, \zeta_\alpha \in \mathbb{F}_p$, the sums that appear are of the form:

$$\sum_{\zeta_1,\dots,\zeta_\alpha\in\mathbb{F}_p} \zeta_1^{i_1} \cdots \zeta_\alpha^{i_\alpha},$$

with $i_1, \dots, i_\alpha \in \mathbb{N}$, and $i_1 + \cdots + i_\alpha \leq r$. Since $r < (p-1)\alpha$, these sums vanish and therefore $S = 0$. □

Let $s \geq 0$ be an integer. For $i = 1, \dots, s$, let $\mathrm{sgn}_i : K_\infty^\times \to \overline{\mathbb{F}}_q^\times$ be a group homomorphism such that $\forall \zeta \in \mathbb{F}_q^\times, \mathrm{sgn}_i(\zeta) = \zeta$. Observe that:

$$\forall i \in \{1, \dots, s\}, \forall x \in K_\infty^\times \cap U_\infty, \quad \mathrm{sgn}_i(x) = 1.$$

For $i = 1, \dots, s$, let $\rho_i : K(\mathrm{sgn}_i(K^\times)) \hookrightarrow F$ be an $\mathbb{F}_p$-algebra homomorphism.

Lemma 3.1 combined with the Riemann-Roch Theorem yields the following result which will be crucial in the sequel.

Lemma 3.2. *Let η be a generator of the cyclic group $\mathbb{F}_\infty^\times$. Let $\mathfrak{I}$ be a non-zero ideal of O_E with $\mathfrak{I} \in \mathcal{I}_{\mathcal{S}}(\mathcal{B})$. Let $\mathbb{F}_E$ be the algebraic closure of $\mathbb{F}_q$ in E. Let k be an integer, $1 \leq k \leq \frac{q^{d_\infty}-1}{q-1}$. Let*

$$S_{m,\mathcal{S}}(\mathfrak{I}, \mathcal{B}, N, k) = \{aO_E \in \mathcal{P}_{\mathcal{S}}(\mathcal{B}, N), \mathrm{sgn}(N_{E/K}(a)) \equiv \eta^k \pmod{\mathbb{F}_q^\times};$$
$$a\mathfrak{I} \subset O_E, \deg(N_{E/K}(a\mathfrak{I})) = m\}.$$

If

$$m \geq 2g_E[\mathbb{F}_E : \mathbb{F}_q] + d_\infty(N) + d_{\mathcal{S}}(\mathcal{B}) + \frac{s[E:K]_{sep}}{(p-1)[\mathbb{F}_q : \mathbb{F}_p]}$$

where $[E:K]_{sep}$ is the separable degree of the finite extension E/K, then we have:

$$\sum_{aO_E \in S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)} \rho_1(\frac{N_{E/K}(a)}{\mathrm{sgn}_1(N_{E/K}(a))}) \cdots \rho_s(\frac{N_{E/K}(a)}{\mathrm{sgn}_s(N_{E/K}(a))}) = 0.$$

In particular, let's set:

$$\begin{aligned} S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N) &= \cup_{k=1}^{\frac{q^{d_\infty}-1}{q-1}} S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k) \\ &= \{aO_E \in \mathcal{P}_{\mathcal{S}}(\mathcal{B},N), a\mathfrak{I} \subset O_E, \deg(N_{E/K}(a\mathfrak{I})) = m\}. \end{aligned}$$

If

$$m \geq 2g_E[\mathbb{F}_E:\mathbb{F}_q] + d_\infty(N) + d_{\mathcal{S}}(\mathcal{B}) + \frac{s[E:K]_{sep}}{(p-1)[\mathbb{F}_q:\mathbb{F}_p]},$$

then we have:

$$\sum_{aO_E \in S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N)} \rho_1(\frac{N_{E/K}(a)}{\mathrm{sgn}_1(N_{E/K}(a))}) \cdots \rho_s(\frac{N_{E/K}(a)}{\mathrm{sgn}_s(N_{E/K}(a))}) = 0.$$

Proof. The proof of the Lemma is based on the arguments used in the proof of [21], Theorem 8.9.2. We give a detailed proof for the convenience of the reader. Since there is a finite number of ideals J of A such that $\deg J = m$, and, given such an ideal J, there is a finite number of ideals $\mathfrak{J}$ in O_E such that $N_{E/K}(\mathfrak{J}) = J$, we deduce that $S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)$ is a finite set, possibly empty. Let's fix $aO_E \in S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)$ (so we assume that $m \geq 0$ is such that $S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k) \neq \emptyset$). Let $V(a)$ be the set of elements $b \in E$ such that

i) $b\mathfrak{I} \subset O_E$,
ii) $v_i(b) \geq v_i(a) + m_i + 1$, for $i = 1, \ldots, r$,
iii) $\mathrm{ord}_{\mathfrak{P}}(b) \geq \mathrm{ord}_{\mathfrak{P}}(\mathcal{B})$, for all $\mathfrak{P}$ dividing $\mathcal{B}$,
iv) $\mathrm{ord}_{\mathfrak{P}}(b) \geq 1$, for all $\mathfrak{P} \in \mathcal{S}$.

Observe that:

$$\deg N_{E/K}(a\mathfrak{I}) = \deg N_{E/K}(\mathfrak{I}) - d_\infty \sum_{i=1}^{r} f_i v_i(a).$$

But:

$$\deg N_{E/K}(\mathfrak{I}) = [\mathbb{F}_E:\mathbb{F}_q] \dim_{\mathbb{F}_E} \frac{O_E}{\mathfrak{I}},$$

therefore:

$$\frac{-1}{[\mathbb{F}_E:\mathbb{F}_q]} \sum_{i=1}^{r} d_\infty f_i v_i(a) = \frac{m}{[\mathbb{F}_E:\mathbb{F}_q]} - \dim_{\mathbb{F}_E} \frac{O_E}{\mathfrak{I}}.$$

We now assume that $m - d_\infty(N) - d_{\mathcal{S}}(\mathcal{B}) \geq [\mathbb{F}_E:\mathbb{F}_q](2g_E - 1)$. Then, by the Riemann-Roch Theorem, $V(a)$ is a finite $\mathbb{F}_E$-vector space of dimension

$$\dim V(a) = \frac{m}{[\mathbb{F}_E:\mathbb{F}_q]} + 1 - g_E - \frac{d_\infty(N) + d_{\mathcal{S}}(\mathcal{B})}{[\mathbb{F}_E:\mathbb{F}_q]}.$$

We have (see [21], Lemma 8.9.3):

$$(a + V(a))O_E \subset S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k),$$
$$\forall b, b' \in V(a), \quad (a+b)O_E = (a+b')O_E \Leftrightarrow b = b'.$$

If $\mathrm{sgn}' : K_\infty^\times \to \overline{\mathbb{F}_q}^\times$ is a group homomorphism, we have:

$$\forall b \in V(a), \quad \mathrm{sgn}'(N_{E/K}(a+b)) = \mathrm{sgn}'(N_{E/K}(a)).$$

Let $cO_E \in S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)$. Then:

$$(c+V(c))O_E \cap (a+V(a))O_E \neq \emptyset \Leftrightarrow cO_E \in (a+V(a))O_E.$$

Thus, if $S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k) \neq \emptyset$, let's select $a_1O_E,\dots,a_lO_E \in S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)$ such that $S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)$ is the disjoint union of the $(a_i+V(a_i))O_E$. We have:

$$\sum_{cO_E \in S_{m,\mathcal{S}}(\mathfrak{I},\mathcal{B},N,k)} \rho_1\Big(\frac{N_{E/K}(c)}{\mathrm{sgn}_1(N_{E/K}(c))}\Big)\cdots\rho_s\Big(\frac{N_{E/K}(c)}{\mathrm{sgn}_s(N_{E/K}(c))}\Big) =$$

$$\sum_{i=1}^{l}\sum_{b\in V(a_i)} \rho_1\Big(\frac{N_{E/K}(a_i+b)}{\mathrm{sgn}_1(N_{E/K}(a_i))}\Big)\cdots\rho_s\Big(\frac{N_{E/K}(a_i+b)}{\mathrm{sgn}_s(N_{E/K}(a_i))}\Big).$$

Let's fix $1 \leq i \leq l$, and let

$$S = \sum_{b\in V(a_i)} \rho_1(N_{E/K}(a_i+b))\cdots\rho_s(N_{E/K}(a_i+b)).$$

Let E_1/K consist of the elements of E which are separable over K. Let $p^{l'} = [E : E_1]$. Then:

$$S = \sum_{b\in V(a_i)} \rho_1(N_{E_1/K}(a_i^{p^{l'}}+b^{p^{l'}}))\cdots\rho_s(N_{E_1/K}(a_i^{p^{l'}}+b^{p^{l'}})).$$

Therefore, we can assume that E/K is a finite separable extension. Let $\overline{F}$ be an algebraic closure of F. Then, for $i=1,\dots,s$, ρ_i extends to a morphism $\rho_i : \overline{K} \to \overline{F}$. Let $\sigma_l : E \to \overline{K}$ be the distincts K-embeddings of E in $\overline{K}$, $l=1,\dots,[E:K]$. Then:

$$\forall i=1,\dots,s, \quad \rho_i(N_{E/K}(a_i+b)) = \prod_{l=1}^{[E:K]} \rho_i(\sigma_l(a_i+b)).$$

By Lemma 3.1, if $[\mathbb{F}_q:\mathbb{F}_p](m+[\mathbb{F}_E:\mathbb{F}_q](1-g_E)-d_\infty(N)-d_{\mathcal{S}}(\mathcal{B})) > \frac{s[E:K]}{p-1}$, we have $S=0$. □

Let $s \geq 0$ be an integer. For $i=1,\dots,s$, let $\psi_i : \mathcal{I}(A) \to \overline{K}_\infty$ be a map such that there exists $M(\psi_i) \subset K_\infty^\times$ an open subgroup of finite index with the following property:

$$\forall I \in \mathcal{I}(A), \forall \alpha \in K^\times \cap M(\psi_i), \quad \psi_i(I\alpha) = \psi_i(I)\frac{\alpha}{\mathrm{sgn}_i(\alpha)}.$$

For $i=1,\dots,s$, let $K(\psi_i) = K(\mathrm{sgn}_i(K^\times), \psi_i(I), I \in \mathcal{I}(A)) \subset \overline{K}_\infty$. Let $\chi : \mathcal{I}(\mathcal{B}) \to F$ be a map such that:

$$\forall \mathfrak{I} \in I(\mathcal{B}), \forall \mathfrak{J} \in \mathcal{P}(\mathcal{B},N), \quad \chi(\mathfrak{I}\mathfrak{J}) = \chi(\mathfrak{I}).$$

For $j=1,\dots,r$, let $N'_j = N_j \cap N^{-1}_{E_{v_j}/K_\infty}(M(\psi_1)\cap\ldots\cap M(\psi_s))$, then N'_j is an open subgroup of finite index of $E_{v_j}^\times$. Let $N' = \prod_{j=1}^r N'_j$.

Corollary 3.3. *Let $s \geq 0$ be an integer and let, for $i=1,\dots,s$, $\rho_i : K(\psi_i) \hookrightarrow F$ be an $\mathbb{F}_p$-algebra homomorphism. If*

$$m \geq 2g_E[\mathbb{F}_E:\mathbb{F}_q] + d_\infty(N') + d_{\mathcal{S}}(\mathcal{B}) + \frac{s[E:K]_{sep}}{(p-1)[\mathbb{F}_q:\mathbb{F}_p]},$$

then:

$$\sum_{\substack{\mathfrak{I}\subset O_E,\mathfrak{I}\in\mathcal{I}_{\mathcal{S}}(\mathcal{B})\\ \deg N_{E/K}(\mathfrak{I})=m}} \chi(\mathfrak{I})\,\rho_1(\psi_1(N_{E/K}(\mathfrak{I})))\cdots\rho_s(\psi_s(N_{E/K}(\mathfrak{I})))=0.$$

Proof. For $m\geq 0$, let's set:

$$U_m=\sum_{\substack{\mathfrak{I}\subset O_E,\mathfrak{I}\in\mathcal{I}_{\mathcal{S}}(\mathcal{B})\\ \deg N_{E/K}(\mathfrak{I})=m}} \chi(\mathfrak{I})\,\rho_1(\psi_1(N_{E/K}(\mathfrak{I})))\cdots\rho_s(\psi_s(N_{E/K}(\mathfrak{I}))).$$

Let $h=\mid \frac{\mathcal{I}(\mathcal{B})}{\mathcal{P}(\mathcal{B},N')} \mid$. Select $\mathfrak{I}_1,\ldots,\mathfrak{I}_h\in\mathcal{I}_{\mathcal{S}}(\mathcal{B})$, $\mathfrak{I}_1,\ldots,\mathfrak{I}_h\subset O_E$, a system of representatives of $\frac{\mathcal{I}(\mathcal{B})}{\mathcal{P}(\mathcal{B},N')}$. Then U_m is equal to

$$\sum_{j=1}^{h}\chi(\mathfrak{I}_j)\prod_{i=1}^{s}\rho_i(\psi_i(N_{E/K}(\mathfrak{I}_j)))\sum_{aO_E\in S_{m,\mathcal{S}}(\mathfrak{I}_j,\mathcal{B},N')}\prod_{i=1}^{s}\rho_i\Big(\frac{N_{E/K}(a)}{\mathrm{sgn}_i(N_{E/K}(a))}\Big).$$

Therefore, by Lemma 3.2, if $m\geq 2g_E[\mathbb{F}_E:\mathbb{F}_q]+d_\infty(N')+d_{\mathcal{S}}(\mathcal{B})+\frac{s[E:K]_{sep}}{(p-1)[\mathbb{F}_q:\mathbb{F}_p]}$, we have:

$$U_m=0.$$

□

Let $s\geq 1$ be an integer. Let $f:\mathbb{Z}\to 1+(q-1)\mathbb{Z}$ be a map. For $i=1,\ldots,s$, let $\psi_i:\mathcal{I}(A)\to\overline{K}_\infty$ be a map such that :

$$\forall I\in\mathcal{I}(A),\forall\alpha\in K^\times,\quad \psi_i(I\alpha)=\psi_i(I)\frac{\alpha}{\mathrm{sgn}_i(\alpha)^{f(\deg I)}}.$$

Let's observe that, by the proof of Lemma 2.1, for $i=1,\ldots,s$, there exist $e_i\in\mathbb{Z}$, $e_i\equiv 1 \pmod{q-1}$, and $\zeta_i\in\overline{\mathbb{F}}_q^\times$ such that:

$$\forall a\in K_\infty^\times,\quad \mathrm{sgn}_i(a)=\mathrm{sgn}(a)^{e_i}\zeta_i^{v_\infty(a)}.$$

Let $\chi:\mathcal{I}(A)\to F$ be a map such that:

$$\forall I\in\mathcal{I}(A),\forall\alpha\in K^\times,\quad \chi(I\alpha)=\chi(I).$$

For $i=1,\ldots,s$, let $\rho_i:K(\psi_i)\hookrightarrow F$ be an $\mathbb{F}_p$-algebra homomorphism. For $i=1,\ldots,s$, let $l_i\in\mathbb{N}$ such that:

$$\forall\alpha\in K^\times,\quad \rho_i(\mathrm{sgn}_i(\alpha))=\mathrm{sgn}_i(\alpha)^{p^{l_i}}.$$

Corollary 3.4. *Let $s\geq 1$. If*

$$p^{l_1}e_1+\cdots+p^{l_s}e_s\equiv 0 \pmod{q^{d_\infty}-1},$$

and

$$\prod_{i=1}^{s}\zeta_i^{p^{l_i}}=1,$$

then :

$$\sum_{m\geq 0}\sum_{\substack{I\in\mathcal{I}(A),I\subset A\\ \deg I=m}}\chi(I)\rho_1(\psi_1(I))\cdots\rho_s(\psi_s(I))=0.$$

Proof. Here $E = K$, $\mathcal{S} = \emptyset$, $\mathcal{B} = A$, $N = K_\infty^\times$. For $m \geq 0$, let's set:

$$U_m = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(A), I \subset A \\ \deg I = m}} \chi(I)\rho_1(\psi_1(I)) \cdots \rho_s(\psi_s(I)).$$

Let $h = \mid \mathrm{Pic}(A) \mid$. Select $I_1, \ldots, I_h \in \mathcal{I}(A)$, $I_1, \ldots, I_h \subset A$, a system of representatives of $\mathrm{Pic}(A)$. Then U_m is equal to:

$$\sum_{j=1}^{h} \chi(I_j) \prod_{i=1}^{s} \rho_i(\psi_i(I_j)) \sum_{aA \in S_{m,\mathcal{S}}(I_j, A, N)} \prod_{i=1}^{s} \rho_i\left(\frac{a}{\mathrm{sgn}_i(a)^{f(\deg I_j)}}\right).$$

Now, assume that $p^{l_1}e_1 + \cdots + p^{l_s}e_s \equiv 0 \pmod{q^{d_\infty} - 1}$, and $\prod_{i=1}^{s} \zeta_i^{p^{l_i}} = 1$. For simplicity, we set:

$$S_m(I_j) = S_{m,\mathcal{S}}(I_j, A, N) = \{aA, a \in K^\times, aI_j \subset A, \deg(aI_j)) = m\}.$$

Then $\sum_{m \geq 0} U_m$ is equal to the finite sum (Lemma 3.2):

$$\sum_{j=1}^{h} \chi(I_j)\rho_1(\psi_1(I_j)) \cdots \rho_s(\psi_s(I_j)) \sum_{m \geq 0} \sum_{aA \in S_m(I_j)} \rho_1(a) \ldots \rho_s(a).$$

Let's select $d \geq 0$ such that:

$$\sum_{m \geq 0} \sum_{aA \in S_m(I_j)} \rho_1(a) \ldots \rho_s(a) = \sum_{m=0}^{d} \sum_{aA \in S_m(I_j)} \rho_1(a) \ldots \rho_s(a).$$

Let $V_d = \{a \in K^\times, aA \in S_m(I_j), m \leq d\} \cup \{0\}$. Then V_d is a finite dimensional $\mathbb{F}_q$-vector space and:

$$\sum_{m=0}^{d} \sum_{aA \in S_m(I_j)} \rho_1(a) \ldots \rho_s(a) = -\sum_{a \in V_d} \rho_1(a) \ldots \rho_s(a).$$

By Lemma 3.1, if $\dim_{\mathbb{F}_q} V_d > \frac{s}{(p-1)[\mathbb{F}_q : \mathbb{F}_p]}$, then:

$$\sum_{a \in V_d} \rho_1(a) \ldots \rho_s(a) = 0.$$

Now, the Riemann-Roch Theorem tells us that, for $d \geq 2g_K - 1$, we have: $\dim_{\mathbb{F}_q} V_d = d + 1 - g_K$. Thus:

$$\sum_{m \geq 0} \sum_{aA \in S_m(I_j)} \rho_1(a) \ldots \rho_s(a) = 0.$$

□

4. Several variable twisted zeta functions

4.1. **The ∞-adic case.**

Let E/K be a finite extension, and let $\mathcal{B}$ and N as in the previous section. Let $\mathcal{S}$ be a finite set, possibly empty, of maximal ideal of O_E which are relatively prime to $\mathcal{B}$. Let $\chi : \mathcal{I}(\mathcal{B}) \to F$ be a map such that:

$$\forall \mathfrak{I} \in \mathcal{I}(\mathcal{B}), \forall \mathfrak{J} \in \mathcal{P}(\mathcal{B}, N), \quad \chi(\mathfrak{I}\mathfrak{J}) = \chi(\mathfrak{I}).$$

Let $s \geq 0, n \geq 1$ be two integers. Let $\psi_1, \ldots, \psi_s, \eta_1, \ldots \eta_n : \mathcal{I}(A) \to \overline{K}_\infty$ be $s+n$ admissible maps. We assume that for $j = 1, \ldots, n$, $\eta_j(\mathcal{I}(A)) \subset \overline{K}_\infty^\times$, and for $i = 1, \ldots, s$, $n(\psi_i) \in \mathbb{N}$. For $i = 1, \ldots, s$, let $\rho_i : K(\psi_i) \hookrightarrow F$ be an $\mathbb{F}_p$-algebra homomorphism. For $i = 1, \ldots, n$, let $\sigma_i : K_\infty(\langle \eta_i \rangle) \hookrightarrow F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. Let $u = (x; z_1, \ldots, z_n) \in \mathbb{S}_{F,n} = F^\times \times \mathbb{Z}_p^n$.

Theorem 4.1. *The following sum converges in F :*

$$\sum_{m \geq 0} \sum_{\substack{\mathfrak{I} \in \mathcal{I}_{\mathcal{S}}(\mathcal{B}), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = m}} \chi(\mathfrak{I}) \rho_1(\psi_1(N_{E/K}(\mathfrak{I}))) \cdots \rho_s(\psi_s(N_{E/K}(\mathfrak{I}))) N_{E/K}(\mathfrak{I})^{-u}_{\underline{\sigma}, \underline{\eta}}.$$

We denote the above sum by $\zeta_{\mathcal{S}, O_E}(\underline{\rho}, \underline{\psi}; \underline{\sigma}, \underline{\eta}; \chi; u)$.

Proof. The proof is in the spirit of the proof of [5], Proposition 6.

Let $\infty_1, \ldots, \infty_r$ be the places of E above ∞. let $v_1, \ldots, v_r$ be the valuation on E associated to these places. We denote by E_{v_i} the v_i-adic completion of E, $i = 1, \ldots, r$. Recall that $N = \prod_{i=1}^r N_i$, where $N_i \subset E_{v_i}^\times$ is an open subgroup of finite index. For $i = 1, \ldots, r$, let:

$$N_i' = N_i \cap N_{E_{v_i}/K_\infty}^{-1}(M(\psi_1) \cap \ldots \cap M(\psi_s) \cap M(\eta_1) \ldots \cap M(\eta_n)).$$

Then $N_i' \subset E_{v_i}^\times$ is also an open subgroup of finite index. Let's set:

$$N' = \prod_{i=1}^r N_i'.$$

Let $\mathfrak{I}_1, \ldots, \mathfrak{I}_h \in \mathcal{I}_{\mathcal{S}}(\mathcal{B})$, $\mathfrak{I}_1, \ldots, \mathfrak{I}_h \subset O_E$ be a system of representatives of $\frac{\mathcal{I}_{\mathcal{S}}(\mathcal{B})}{\mathcal{P}_{\mathcal{S}}(\mathcal{B}, N')}$. Recall that

$$S_{m,\mathcal{S}}(\mathfrak{I}_j, \mathcal{B}, N') = \{aO_E \in \mathcal{P}_{\mathcal{S}}(\mathcal{B}, N'), a\mathfrak{I}_j \subset O_E, \deg(N_{E/K}(a\mathfrak{I}_j)) = m\}.$$

Let's define $V_{m,j}$ to be the following sum:

$$\sum_{aO_E \in S_{m,\mathcal{S}}(\mathfrak{I}_j, \mathcal{B}, N')} (\prod_{i=1}^s \rho_i(\gamma_{\psi_i} \pi^{-n(\psi_i)}))^{v_\infty(N_{E/K}(a))} \prod_{i=1}^s \rho_i(\frac{N_{E/K}(a)}{\operatorname{sgn}(N_{E/K}(a))})^{n(\psi_i)} \times$$
$$\times (\prod_{j=1}^n \sigma_j(\langle \gamma_{\eta_j} \rangle^{-z_j}))^{v_\infty(N_{E/K}(a))} \prod_{j=1}^n \sigma_j(\langle N_{E/K}(a) \rangle)^{-z_j n(\eta_j)}.$$

Then:

$$x^m \sum_{\substack{\mathfrak{I} \in \mathcal{I}_{\mathcal{S}}(\mathcal{B}), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = m}} \chi(\mathfrak{I}) \rho_1(\psi_1(N_{E/K}(\mathfrak{I}))) \cdots \rho_s(\psi_s(N_{E/K}(\mathfrak{I}))) N_{E/K}(\mathfrak{I})^{-u}_{\underline{\sigma}, \underline{\eta}}$$
$$= \sum_{j=1}^h \chi(\mathfrak{I}_j) \rho_1(\psi_1(N_{E/K}(\mathfrak{I}_j))) \cdots \rho_s(\psi_s(N_{E/K}(\mathfrak{I}_j))) N_{E/K}(\mathfrak{I}_j)^{-\underline{z}}_{\underline{\sigma}, \underline{\eta}} V_{m,j}.$$

Note that $v_F \circ \rho_i \mid_K$ is a valuation on K, thus, it is equivalent to the trivial valuation or to the valuation attached to a place of K. This implies that there exists integers $D, D' \in \mathbb{R}$, such that for all $m \geq 0$, for all $j \in \{1, \dots, h\}$ and for all $aO_E \in S_{m,\mathcal{S}}(\mathfrak{I}_j, \mathcal{B}, N')$, we have:

$$v_F(\prod_{i=1}^{s} \rho_i(N_{E/K}(a))^{n(\psi_i)}) \geq -Dm - D'.$$

If $l \in \mathbb{N}$, we denote by $\ell_p(l)$ the sum of the digits of l in base p. By Lemma 3.2, if $m_1, \dots m_n \in \mathbb{N}$ are such that $m_1 + \ldots + m_n \geq 1$ and if

$$m \geq 2g_E[\mathbb{F}_E : \mathbb{F}_q] + d_\infty(N') + d_{\mathcal{S}}(\mathcal{B}) + \frac{(s + \sum_{i=1}^{n} \ell_p(m_i))[E:K]_{sep}}{(p-1)[\mathbb{F}_q : \mathbb{F}_p]},$$

then:

$$\sum_{aO_E \in S_{m,\mathcal{S}}(\mathfrak{I}_j, \mathcal{B}, N')} \prod_{i=1}^{s} \rho_i(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))}) \prod_{j=1}^{n} \sigma_j(\langle N_{E/K}(a) \rangle)^{m_j} = 0.$$

Now, let $l \in \mathbb{N}$, and select $k_1, \dots, k_n \in \mathbb{N} \setminus \{0\}$ such that:
i) $\ell_p(k_i) \leq (l+1)(p-1)$,
ii) $z_i n(\eta_i) + k_i \equiv 0 \pmod{p^{l+1}\mathbb{Z}_p}$.
For example, write $-z_i = \sum_{j \geq 0} u_{i,j} p^j$, $u_{i,j} \in \{0, \dots, p-1\}$, and set $k_i = \sum_{j=0}^{l} u_{i,j} p^j + p^{l+1}$. Let

$$C = [E:K]_{sep}(p-1)[\mathbb{F}_q : \mathbb{F}_p](2g_E[\mathbb{F}_E : \mathbb{F}_q] + d_\infty(N') + d_{\mathcal{S}}(\mathcal{B}) + (\sum_{i=1}^{s} n(\psi_i)) + 1).$$

Then, if $m \geq Cn(l+1)$, we have:

$$\sum_{aO_E \in S_{m,\mathcal{S}}(\mathfrak{I}_j, \mathcal{B}, N')} \prod_{i=1}^{s} \rho_i(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})^{n(\psi_i)} \prod_{j=1}^{n} \sigma_j(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})^{k_j} = 0.$$

Thus, there exists $D'' \in \mathbb{R}$ such that:

$$v_F(V_{m,j}) \geq p^{l+1}\mathrm{Inf}\{v_F(y_i), i = 1, \dots, n\} - D''m - D',$$

where $y_i = \sigma_i(\pi), i = 1, \dots, n$. Note also that $v_F(y_i) > 0, i = 1, \dots, n$. Therefore, if $m \geq Cn$, we get:

$$v_F(V_{m,j}) \geq p^{[\frac{m}{Cn}]}\mathrm{Inf}\{v_F(y_i), i = 1, \dots, n\} - D''m - D'.$$

Therefore, for $m \geq Cn$, there exists $D''' \in \mathbb{R}$ such that we have:

$$v_F\Big(x^m \sum_{\substack{\mathfrak{I} \in \mathcal{I}_{\mathcal{S}}(\mathcal{B}), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = m}} \chi(\mathfrak{I}) \rho_1(\psi_1(N_{E/K}(\mathfrak{I}))) \cdots \rho_s(\psi_s(N_{E/K}(\mathfrak{I}))) N_{E/K}(\mathfrak{I})^{-u}_{\underline{\sigma},\underline{\eta}}\Big)$$
$$\geq p^{[\frac{m}{Cn}]}\mathrm{Inf}\{v_F(y_i), i = 1, \dots, n\} - D''m - D'''.$$

□

Remark 4.2. *By the proof of Theorem 4.1, the valuation v_F of the coefficient of x^{-m} in $\zeta_{\mathcal{S},O_E}(\underline{\rho}, \underline{\psi}; \underline{\sigma}, \underline{\eta}; \chi; u)$ grows exponentially in m for large m.*

Assume that $\psi_1, \dots, \psi_s, \eta_1, \dots \eta_n$, and χ are group homomorphisms, then note that if $u = (x; z_1, \dots, z_n)$ with $v_F(x) << 0$, we have:

$$\zeta_{\mathcal{S},O_E}(\underline{\rho}, \underline{\psi}; \underline{\sigma}, \underline{\eta}; \chi; u) = \prod_{\mathfrak{P}} (1 - \chi(\mathfrak{P})\rho_1(\psi_1(N_{E/K}(\mathfrak{P}))) \cdots \rho_s(\psi_s(N_{E/K}(\mathfrak{P})))N_{E/K}(\mathfrak{P})^{-u}_{\underline{\sigma},\underline{\eta}})^{-1},$$

where $\mathfrak{P}$ runs through the set of maximal ideals of O_E that are contained in $\mathcal{I}_{\mathcal{S}}(\mathcal{B})$.

Let $f : \mathbb{Z} \to 1 + (q-1)\mathbb{Z}$ be a map. For $i = 1, \dots, s$, let $\mathrm{sgn}_i : K_\infty^\times \to \overline{\mathbb{F}}_q^\times$ be a group homomorphism such that $\mathrm{sgn}_i \mid_{\mathbb{F}_q^\times} = \mathrm{Id}_{\mathbb{F}_q^\times}$. For $i = 1, \dots, s$, let $\psi_i : \mathcal{I}(A) \to \overline{K}_\infty$ be a map such that:

$$\forall I \in \mathcal{I}(A), \forall \alpha \in K^\times, \quad \psi_i(I\alpha) = \psi_i(I)\frac{\alpha}{\mathrm{sgn}_i(\alpha)^{f(\deg I)}}.$$

For $i = 1, \dots, s$, let $\rho_i : K(\mathrm{sgn}_i(K^\times), \psi_i(I), I \in \mathcal{I}(A)) \to F$ be an $\mathbb{F}_p$-algebra homomorphism. For $j = 1, \dots, n$, let $f_j : \mathbb{Z} \to 1 + (q-1)\mathbb{Z}$ be a map, and let $\eta_j : \mathcal{I}(A) \to \overline{K}_\infty^\times$ be a map such that

$$\forall I \in \mathcal{I}(A), \forall \alpha \in K^\times, \quad \eta_j(I\alpha) = \eta_j(I)\frac{\alpha}{\mathrm{sgn}'_j(\alpha)^{f_j(\deg I)}},$$

where $\mathrm{sgn}'_j : K_\infty^\times \to \overline{\mathbb{F}}_q^\times$ is a group homomorphism such that $\mathrm{sgn}'_j \mid_{\mathbb{F}_q^\times} = \mathrm{Id}_{\mathbb{F}_q^\times}$. For $j = 1, \dots, n$, let $\sigma_j : K_\infty(\mathrm{sgn}'_j(K^\times), \pi^{\frac{1}{d_\infty}}, \langle \eta_j(I) \rangle, I \in \mathcal{I}(A)) \to F$ be a continuous $\mathbb{F}_p$-algebra homomorphism.

Let's define $l_1, \dots, l_s, l'_1, \dots, l'_n \in \mathbb{N}$ as follows:

$$\forall i = 1, \dots, s, \forall \alpha \in K^\times, \quad \rho_i(\mathrm{sgn}_i(\alpha)) = \mathrm{sgn}_i(\alpha)^{p^{l_i}},$$
$$\forall j = 1, \dots, n, \forall \zeta \in \mathbb{F}_\infty^\times, \quad \sigma_j(\zeta) = \zeta^{p^{l'_j}}.$$

For $i = 1, \dots, s$, there exist $e_i \in \mathbb{Z}$, $e_i \equiv 1 \pmod{q-1}$, and $\zeta_i \in \overline{\mathbb{F}}_q^\times$ such that:

$$\forall a \in K_\infty^\times, \quad \mathrm{sgn}_i(a) = \mathrm{sgn}(a)^{e_i}\zeta_i^{v_\infty(a)}.$$

We assume that:

$$\forall a \in K^\times, \quad \prod_{i=1}^{s} (\frac{\mathrm{sgn}_i(a)}{\mathrm{sgn}(a)})^{e_i p^{l_i}} = 1, \text{ and } \quad \prod_{i=1}^{s} \zeta_i^{p^{l_i}} = 1.$$

Let $\chi : \mathcal{I}(A) \to F$ be a map such that:

$$\forall I \in \mathcal{I}(A), \forall \alpha \in K^\times, \quad \chi(I\alpha) = \chi(I).$$

Note that $\psi_1, \dots, \psi_s, \eta_1, \dots, \eta_n$ are admissible maps.

Corollary 4.3. *We assume that $E = K$, $\mathcal{B} = A$, $N = K_\infty^\times$, $\mathcal{S} = \emptyset$. Let $m_1, \dots, m_n \in \mathbb{N}$ such that $p^{l_1}e_1 + \cdots + p^{l_s}e_s + m_1 p^{l'_1} + \ldots + m_n p^{l'_n} \equiv 0 \pmod{q^{d_\infty} - 1}$, and $p^{l_1}e_1 + \cdots + p^{l_s}e_s + m_1 p^{l'_1} + \ldots + m_n p^{l'_n} \geq 1$. Then:*

$$\zeta_{\mathcal{S},A}(\underline{\rho}, \underline{\psi}; \underline{\sigma}, \underline{\eta}; \chi; (\sigma_1(\pi^{\frac{1}{d_\infty}})^{m_1} \cdots \sigma_n(\pi^{\frac{1}{d_\infty}})^{m_n}; -m_1, \dots, -m_n)) = 0.$$

Proof. For $i = 1, \dots, n$, let $y_i = \sigma_i(\pi^{\frac{1}{d_\infty}})$. Let $u = (y_1^{m_1} \cdots y_n^{m_n}; -m_1, \dots, -m_n) \in \mathbb{S}_{F,n}$. Let $h = \mid \mathrm{Pic}(A) \mid$. Select $I_1, \dots, I_h \in \mathcal{I}(A)$, $I_1, \dots, I_h \subset A$, a system of

representatives of $\mathrm{Pic}(A)$. Let's set:

$$V_{m,j} = \sum_{aA \in S_{m,\mathcal{S}}(I_j,A,N)} \prod_{i=1}^{s} \rho_i\Big(\frac{a}{\mathrm{sgn}_i(a)^{f(\deg I_j)}}\Big)(aA)^{(m_1,\dots,m_n)}_{\underline{\sigma},\underline{\eta}}.$$

Then, we have:

$$V_{m,j} = \sum_{aA \in S_{m,\mathcal{S}}(I_j,A,N)} \prod_{i=1}^{s} \frac{\rho_i(a)}{\mathrm{sgn}(a)^{p^{l_i}e_i}} \prod_{j=1}^{n} \sigma_j\Big(\frac{a}{\mathrm{sgn}(a)\pi^{v_\infty(a)}}\Big)^{m_j}.$$

Since $p^{l_1}q^{e_1} + \cdots + p^{l_s}q^{e_s} + m_1 p^{l'_1} + \ldots + m_n p^{l'_n} \equiv 0 \pmod{q^{d_\infty} - 1}$ and $x = y_1^{m_1} \cdots y_n^{m_n}$,

$$V_{m,j}x^{-m} = -\Big(\prod_{j=1}^{n} y_j^{-m_j \deg I_j}\Big) \sum_{\substack{a \in K^\times \\ aA \in S_{m,\mathcal{S}}(I_j,A,N)}} \rho_1(a)\dots\rho_s(a)\sigma_1(a)^{m_1}\cdots\sigma_n(a)^{m_n}.$$

The assertion is then a consequence of the proof of Corollary 3.4. □

4.2. **The case of finite places.**

Let v be a finite place of K, i.e. $v : K \to \mathbb{Z} \cup \{+\infty\}$ is a discrete valuation on K such that there exists $a \in A$ with $v(a) = 1$. Let $P_v = \{a \in A, v(a) > 0\}$ be the maximal ideal of A corresponding to the finite place v. We denote by K_v the v-adic completion of K. We fix $\overline{K}_v$ an algebraic closure of K_v. Then v extends to a valuation $v : \overline{K}_v \to \mathbb{Q} \cup \{+\infty\}$. We fix a K-morphism $\overline{K} \hookrightarrow \overline{K}_v$, and, by abuse of notation, we will identify the elements in $\overline{K}$ with their images in $\overline{K}_v$. We fix π_v a prime of K_v, i.e. $\pi_v \in K_v$ and $v(\pi_v) = 1$. Then, as in section 2, $\forall x \in \overline{K}_v^\times$, x can be written in a unique way:

$$x = \pi_v^{v(x)}\omega_v(x)\langle x\rangle_v \text{ with } \quad \omega_v(x) \in \overline{\mathbb{F}}_q^\times, v(\langle x\rangle_v - 1) > 0.$$

Let $\psi_1, \dots, \psi_s, \eta_1, \dots \eta_n$ be $s+n$ admissible maps. We assume that $n(\psi_i) \in \mathbb{N}, i = 1, \dots, s$, and $n(\eta_j) \in \mathbb{Z}, j = 1, \dots, n$, and that our $s+n$ admissible maps are algebraic. We assume that, for $j = 1, \dots, n$, $\eta_j(\mathcal{I}(A)) \subset \overline{K}_v^\times$. For $i = 1, \dots, s$, we recall that $K(\psi_i) = K(\mathbb{F}_\infty, \gamma_{\psi_i}\pi^{-n(\psi_i)}, \psi_i(I), I \in \mathcal{I}(A)) \subset \overline{K}_v$. For $j = 1, \dots, n$, we set $K_v(\langle\eta_j\rangle_v) := K_v(\langle\eta_j(I)\rangle_v, I \in \mathcal{I}(A)) \subset \overline{K}_v$. Observe that, for $j = 1, \dots, n$, $K_v(\langle\eta_j\rangle_v)/K_v$ is a finite extension.

Let (F, v_F) be a field which is complete with respect to a non-trivial valuation $v_F : F \to R \cup \{+\infty\}$ and such that F is an $\overline{\mathbb{F}}_q$-algebra.

Let $n \geq 1$, and let's set:

$$\mathbb{S}_{v,F,n} = F^\times \times \mathbb{Z}_p^n \times \mathbb{Z}^n.$$

For $i = 1, \dots, n$, let $\sigma_i : K_v(\langle\eta_i\rangle_v) \hookrightarrow F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. For $u = (x; z_1, \dots, z_n; \underline{\delta}) \in \mathbb{S}_{v,F,n}$, $I \in \mathcal{I}(A)$, we set:

$$I^u_{\underline{\sigma},\underline{\eta}} = x^{\deg I}\omega_v(\eta_1(I))^{\delta_1}\cdots\omega_v(\eta_n(I))^{\delta_n}\sigma_1(\langle\eta_1(I)\rangle_v^{z_1})\cdots\sigma_n(\langle\eta_n(I)\rangle_v^{z_n}) \in F^\times.$$

Let E/K be a finite extension, and let $\mathcal{B}$ and N as in the previous section. Let $\mathcal{S}$ be a finite set, possibly empty, of maximal ideals of O_E which are relatively prime to $\mathcal{B}$. Let $\mathcal{S}_v$ be the union of S and the maximal ideals of O_E above P_v and that do not divide $\mathcal{B}$. Let $\chi : \mathcal{I}(\mathcal{B}) \to F$ be a map such that:

$$\forall \mathfrak{I} \in \mathcal{I}(\mathcal{B}), \forall \mathfrak{J} \in \mathcal{P}(\mathcal{B}, N), \quad \chi(\mathfrak{I}\mathfrak{J}) = \chi(\mathfrak{I}).$$

For $i = 1, \dots, s$, let $\rho_i : K(\psi_i) \hookrightarrow F$ be an $\mathbb{F}_p$-algebra homomorphism. Let $u = (x; z_1, \dots, z_n; \underline{\delta}) \in \mathbb{S}_{v,F,n}$.

Theorem 4.4. *The following sum converges in F :*

$$\sum_{m\geq 0} \sum_{\substack{\mathfrak{I}\in\mathcal{I}_{\mathcal{S}_v}(\mathcal{B}),\mathfrak{I}\subset O_E \\ \deg(N_{E/K}(\mathfrak{I}))=m}} \chi(\mathfrak{I})\rho_1(\psi_1(N_{E/K}(I)))\cdots\rho_s(\psi_s(N_{E/K}(I)))N_{E/K}(\mathfrak{I})^{-u}_{\underline{\sigma},\underline{\eta}}.$$

Proof. The proof of the Theorem is similar to the proof of Theorem 4.1. We only give a sketch of the proof. Let N' be as in the proof of Theorem 4.1.

Let $\mathfrak{I}_1, \dots, \mathfrak{I}_h \in \mathcal{I}_{\mathcal{S}_v}(\mathcal{B})$, $\mathfrak{I}_1, \dots, \mathfrak{I}_h \subset O_E$ be a system of representatives of $\frac{\mathcal{I}(\mathcal{B})}{\mathcal{P}(\mathcal{B},N')}$. Recall that

$$S_{m,\mathcal{S}_v}(\mathfrak{I}_j, \mathcal{B}, N') = \{aO_E \in \mathcal{P}_{\mathcal{S}_v}(\mathcal{B}, N'), a\mathfrak{I}_j \subset O_E, \deg(N_{E/K}(a\mathfrak{I}_j)) = m\}.$$

Let $V_{m,j}$ be the following sum:

$$\sum_{aO_E\in S_{m,\mathcal{S}_v}(\mathfrak{I}_j,\mathcal{B},N')} (\prod_{i=1}^{s}\rho_i(\gamma_{\psi_i}\pi^{-n(\psi_i)}))^{v_\infty(N_{E/K}(a))}\omega_v(\prod_{j=1}^{n}\sigma_j(\gamma_{\psi_j}\pi^{-n(\eta_j)})^{-\delta_j})^{v_\infty(N_{E/K}(a))})$$
$$\times (\prod_{j=1}^{n}\langle\sigma_j(\gamma_{\psi_j}\pi^{-n(\eta_j)})\rangle_v^{-z_j})^{v_\infty(N_{E/K}(a))}\,\omega_v(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})^{-\delta_1 n(\eta_1)-\cdots-\delta_n n(\eta_n)}$$
$$\times \prod_{i=1}^{s}\rho_i(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})^{n(\psi_i)}\prod_{j=1}^{n}\langle\sigma_j(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})\rangle_v^{-z_j n(\eta_j)}.$$

Note that there exist integers $D, D' \in \mathbb{R}$, such that, for $m \geq 0$, $j \in \{1, \dots, r\}$ and $aO_E \in S_{m,\mathcal{S}_v}(\mathfrak{I}_j, \mathcal{B}, N')$, we have:

$$v_F(\prod_{i=1}^{s}\rho_i(N_{E/K}(a))^{n(\psi_i)}) \geq -Dm - D'.$$

By the proof of Lemma 3.2, there exists an integer $C \geq 1$ such that, for $m_1, \dots m_n \in \mathbb{N}$, with $m_1 + \ldots + m_n \geq 1$, $m \geq \frac{C(\sum_{i=1}^{n}\ell_p(m_i))}{p-1}$ and $\delta \in \mathbb{Z}$, we have:

$$\sum_{aO_E\in S_{m,\mathcal{S}_v}(\mathfrak{I}_j,\mathcal{B},N')} \omega_v(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})^{-\delta}\prod_{i=1}^{s}\rho_i(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})^{n(\psi_i)}\times$$
$$\times\prod_{j=1}^{n}\sigma_j\Big(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))\omega_v(\frac{N_{E/K}(a)}{\mathrm{sgn}(N_{E/K}(a))})}\Big)^{m_j} = 0.$$

Now, let $l \in \mathbb{N}$, and select $k_1, \dots, k_n \in \mathbb{N} \setminus \{0\}$ such that:
i) $\ell_p(k_i) \leq (l+1)(p-1)$,
ii) $z_i\eta(\eta_i) + k_i \equiv 0 \pmod{p^{l+1}\mathbb{Z}_p}$.
Then, there exists $D'' \in \mathbb{R}$, such that if $m \geq Cn(l+1)$, we have:

$$v_F(V_{m,j}) \geq p^{l+1}\mathrm{Inf}\{v_F(y_i), i = 1, \dots, n\} - D''m - D',$$

where $y_i = \sigma_i(\pi_v)$. We conclude as in the proof of Theorem 4.1. □

5. Examples

5.1. **The case $A = \mathbb{F}_q[\theta]$.**

We take $\pi = \theta^{-1}$. In that case $d_\infty = 1$, $\mathrm{Pic}(A) = \{1\}$, and we set

$$A_+ = \{a \in A, a \,\text{monic}\} = \{a \in A, \mathrm{sgn}(a) = 1\}.$$

Let $\varphi : \mathbb{F}_q \to \mathbb{F}_q, x \mapsto x^p$. Let $a = \sum_{i=0}^m \lambda_i \theta^i, \lambda_i \in \mathbb{F}_q$, we set:

$$\forall j \geq 0, \quad \varphi^j(a) = \sum_{i=0}^m \lambda_i^{p^j} \theta^i.$$

Let $s \geq 0$, $n \geq 1$, and $e_1, \ldots, e_s, l_1, \ldots, l_n \in \mathbb{N}$. Let (F, v_F) be a complete field with respect to a non-trivial valuation v_F and such that F is an $\mathbb{F}_q$-algebra. Recall that $\mathbb{S}_{F,n} = F^\times \times \mathbb{Z}_p^n$. Let E/K be a finite extension. Let E_1/E be a finite abelian extension and let $(., E_1/E)$ be the Artin symbol. Let $\chi : \mathrm{Gal}(E_1/E) \to F^\times$ be a group homomorphism. If $\mathfrak{P}$ is a maximal ideal of O_E, we set:

$$\chi(\mathfrak{P}) = \begin{cases} 0 & \text{if } \mathfrak{P} \text{ is ramified in } E_1/E, \\ \chi((\mathfrak{P}, E_1/E)) & \text{otherwise.} \end{cases}$$

If we apply Theorem 4.1, we get:

Corollary 5.1. *Let $x_1, \ldots, x_s \in F$, and let $y_1, \ldots, y_n \in F$ such that $v_F(y_i) < 0, i = 1, \ldots, n$. Then, for $z_1, \ldots, z_n \in \mathbb{Z}_p$ and for $x \in F$, the following sum converges in F :*

$$\sum_{m \geq 0} \Big(\sum_{\substack{\mathfrak{I} \in \mathcal{I}(O_E), \mathfrak{I} \subset O_E, \\ N_{E/K}(\mathfrak{I}) = aA, \\ a \in A_+, \deg_\theta a = m}} \chi(\mathfrak{I}) \prod_{i=1}^s \varphi^{e_i}(a) \mid_{\theta = x_i} \prod_{j=1}^n \langle \varphi^{l_j}(a) \mid_{\theta = y_j} \rangle^{z_j} \Big) x^m,$$

where for $y \in F, v_\infty(y) < 0$, and for $a \in A_+$, $\langle a(y) \rangle = \frac{a(y)}{y^{\deg_\theta a}}$.

Let X is an indeterminate over K, and write:

$$\forall a \in A, \quad a(\theta + X) = \sum_{m \geq 0} a^{(m)} X^m, a^{(m)} \in A.$$

Then, for all $m \geq 0$, $.^{(m)} : A \to A$ is an $\mathbb{F}_q$-linear map and we have:

$$\forall k \geq 0, \quad (\theta^k)^{(m)} = \binom{k}{m} \theta^{k-m},$$

where

$$\binom{k}{m} = \begin{cases} 0 & \text{if } k < m, \\ \frac{k!}{m!(k-m)!} \pmod p & \text{if } k \geq m. \end{cases}$$

Observe that:

$$\forall i \geq 0, \forall m \geq 0, \forall a \in A, \quad \varphi^i(a^{(m)}) = (\varphi^i(a))^{(m)}.$$

Proposition 5.2. *Let $x_1, \ldots, x_s \in F$, $m_1, \ldots, m_s \in \mathbb{N}$, and let $y_1, \ldots, y_n \in F$ such that $v_F(y_i) < 0, i = 1, \ldots, n$. Then, for $z_1, \ldots, z_n \in \mathbb{Z}_p$ and for $x \in F$, the following sum converges in F :*

$$S(m_1, \ldots, m_s) := \sum_{m \geq 0} \Big(\sum_{\substack{\mathfrak{I} \in \mathcal{I}(O_E), \mathfrak{I} \subset O_E, \\ N_{E/K}(\mathfrak{I}) = aA, \\ a \in A_+, \deg_\theta a = m}} \chi(\mathfrak{I}) \prod_{i=1}^s \varphi^{e_i}(a^{(m_i)}) |_{\theta = x_i} \prod_{j=1}^n \langle \varphi^{l_j}(a) |_{\theta = y_j} \rangle^{z_j} \Big) x^m.$$

Furthermore:

$$\lim_{m_1+\ldots+m_s\to+\infty} S(m_1,\ldots,m_s)=0.$$

Proof. Let $t_1,\ldots,t_s$ be s indeterminates over F and let $\mathbb{T}_s(F)$ be the Tate algebra in the variables $t_1,\ldots,t_s$ with coefficients in F. Let F' be the completion of the field of fraction of $\mathbb{T}_s(F)$. Let $x_1,\ldots,x_s\in F$, $y_1,\ldots,y_n\in F$ such that $v_F(y_i)<0, i=1,\ldots,n$, $z_1,\ldots,z_n\in\mathbb{Z}_p$, and $x\in F$. By Corollary 5.1, the following sum converges in F' :

$$S:=\sum_{m\geq 0}\Big(\sum_{\substack{\mathfrak{I}\in\mathcal{I}(O_E),\mathfrak{I}\subset O_E,\\ N_{E/K}(\mathfrak{I})=aA,\\ a\in A_+,\deg_\theta a=m}}\chi(\mathfrak{I})\prod_{i=1}^{s}\varphi^{e_i}(a)\mid_{\theta=t_i+x_i}\prod_{j=1}^{n}\langle\varphi^{l_j}(a)\mid_{\theta=y_j}\rangle^{z_j}\Big)x^m.$$

Since $\mathbb{T}_s(F)$ is closed in F', we get $S\in\mathbb{T}_s(F)$. For $i=1,\ldots,s$, we have:

$$\varphi^{e_i}(a)\mid_{\theta=t_i+x_i}=\sum_{m\geq 0}\varphi^{e_i}(a^{(m)})\mid_{\theta=x_i}t_i^m.$$

But, we have:

$$S=\sum_{m_1,\ldots,m_s\in\mathbb{N}}S(m_1,\ldots,m_s)t_1^{m_1}\cdots t_s^{m_s},$$

where $S(m_1,\ldots,m_s)\in F$, and $\lim_{m_1+\ldots+m_s\to+\infty}S(m_1,\ldots,m_s)=0$. Therefore, we get:

$$S(m_1,\ldots,m_s)=\sum_{m\geq 0}\Big(\sum_{\substack{\mathfrak{I}\in\mathcal{I}(O_E),\mathfrak{I}\subset O_E,\\ N_{E/K}(\mathfrak{I})=aA,\\ a\in A_+,\deg_\theta a=m}}\chi(\mathfrak{I})\prod_{i=1}^{s}\varphi^{e_i}(a^{(m_i)})|_{\theta=x_i}\prod_{j=1}^{n}\langle\varphi^{l_j}(a)|_{\theta=y_j}\rangle^{z_j}\Big)x^m.$$

□

We refer the interested reader to [5] and [11] for the arithmetic properties of a special case of the above sums.

Recall that $\mathbb{C}_\infty$ be the completion of $\overline{K}_\infty$. Let $t_1,\ldots,t_s,z$ be $s+1$ indeterminates over $\mathbb{C}_\infty$, and let $\mathbb{T}$ be the Tate algebra in the variables $t_1,\ldots,t_s,z$ with coefficients in $\mathbb{C}_\infty$. Let F be the completion of the field of fractions of $\mathbb{T}$. Take in Corollary 5.1 $E_1=E=K$, $e_1=\ldots=e_s=l_1=\ldots=l_n=0$, $n=1$, $x_i=t_i$, $x=z$, $y_1=\theta$, $z_1=y\in\mathbb{Z}_p$, we get that the following infinite sum converges in $\mathbb{T}$ and is in fact an entire function on $\mathbb{C}_\infty^{s+1}$:

$$\sum_{m\geq 0}\Big(\sum_{a\in A_+,\deg_\theta a=m}\frac{a(t_1)\cdots a(t_s)}{\langle a\rangle^y}\Big)z^m.$$

The above sums were introduced in [28]. In particular, for all $n\in\mathbb{Z}$, the sum

$$L(n;\underline{t};z):=\sum_{m\geq 0}\Big(\sum_{a\in A_+,\deg_\theta a=m}\frac{a(t_1)\cdots a(t_s)}{a^n}\Big)z^m\in\mathbb{T}$$

defines an entire function on $\mathbb{C}_\infty^{s+1}$. Observe that, by Corollary 3.3 and Corollary 3.4, if $n\leq 0$, this sum is finite and furthermore it vanishes at $z=1$ if $s-n\equiv 0\pmod{q-1}$, $s-n\geq 1$. Using a special case of Anderson's log-algebraicity Theorem for $\mathbb{F}_q[\theta]$ ([1], [2], [38] paragraphs 8.9 and 8.10, see also [9], [8], [10] and

the forthcoming work of M. Papanikolas [27]), F. Pellarin proved ([28]), for $s = 1$, a formula connecting $L(1; t_1; 1)$ to a special function introduced by G. Anderson and D. Thakur ([3]). This formula reflects an analytic class number formula à la Taelman ([35], [36], [17], [18]) for $L(1; \underline{t}; 1)$ (see [8]). Such an analytic class number formula has been generalized in [16] to a larger class of L-series (in particular for $L(n; \underline{t}; z)$, for $n \geq 1$). We also refer the reader to [5], [6], [23], [30], [31], [32], [33], [34], [37] for various arithmetic and analytic properties of the series $L(n; \underline{t}; z), n \in \mathbb{Z}$.

Now, let P be a monic irreducible polynomial in A of degree d. Let $\mathbb{C}_P$ be the completion of an algebraic closure K_P of the P-adic completion of K. Let A_P be the valuation ring of K_P. Then:

$$\forall a \in A_P^\times, \quad a = \omega_P(a)\langle a\rangle_P, \text{ with } v_P(\langle a\rangle_P - 1) \geq 1, \omega_P(a) \in \mathbb{F}_{q^d}^\times.$$

Note that $\frac{1}{q^d-1} \in \mathbb{Z}_p^\times$, thus:

$$\forall a \in A_P^\times, \quad \langle a\rangle_P = (a^{q^d-1})^{\frac{1}{q^d-1}}.$$

Let (F, v_F) be a complete field with respect to a non-trivial valuation v_F and such that F is an $\mathbb{F}_{q^d}$-algebra. Let $\sigma : K_P \hookrightarrow F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. Then:

$$\forall a \in A_+, \quad \sigma(\langle a\rangle_P) = (\sigma(a)^{q^d-1})^{\frac{1}{q^d-1}}.$$

Let $y = \sigma(\theta)$, then $v_F(y) \geq 0$, and $y \notin \mathbb{F}_{q^d}$. Furthermore, there exists $i \geq 0$, such that:

$$\forall a \in A, \quad \sigma(a) = \varphi^i(a) \mid_{\theta=y}.$$

Thus:

$$v_F(\varphi^i(P) \mid_{\theta=y}) > 0.$$

Furthermore:

$$\sigma(\langle a\rangle_P) = ((\varphi^i(a) \mid_{\theta=y})^{q^d-1})^{\frac{1}{q^d-1}} =: \langle \varphi^i(a) \mid_{\theta=y}\rangle_P.$$

Let E/K be a finite extension and let E_1/E be a finite abelian extension. Let $\chi : \mathrm{Gal}(E_1/E) \to F^\times$ be a group homomorphism. Let $\mathcal{S}_P$ be the set of maximal ideals of O_E above P. If we apply Theorem 4.4, then, by the proof of Proposition 5.2, we get:

Corollary 5.3. *Let $e_1, \ldots, e_s, m_1, \ldots, m_s, l_1, \ldots, l_n \in \mathbb{N}$. Let $x_1, \ldots, x_s \in F$. Let $y_1, \ldots, y_n \in F \setminus \mathbb{F}_{q^d}$ such that $v_F(\varphi^{l_i}(P) \mid_{\theta=y_i}) > 0$. Then for $z_1, \ldots, z_n \in \mathbb{Z}_p$, for $\delta \in \mathbb{Z}$, and for $x \in F$, the following sum converges in F :*

$$\sum_{m\geq 0}\Big(\sum_{\substack{\mathfrak{I}\in\mathcal{I}_{\mathcal{S}_P}(O_E),\mathfrak{I}\subset O_E,\\ N_{E/K}(\mathfrak{I})=aA,\\ a\in A_+,\deg_\theta a=m}} \omega_P(a)^\delta \chi(\mathfrak{I}) \prod_{i=1}^{s} \varphi^{e_i}(a^{(m_i)})|_{\theta=x_i} \prod_{j=1}^{n} \langle \varphi^{l_j}(a)|_{\theta=y_j}\rangle_P^{z_j}\Big) x^m.$$

Let $t_1, \ldots, t_s, z$ be $s+1$ indeterminates over $\mathbb{C}_P$, and let $\mathbb{T}_P$ be the Tate algebra in the variables $t_1, \ldots, t_s, z$ with coefficients in $\mathbb{C}_P$. Let F_P be the completion of the field of fractions of $\mathbb{T}_P$. Take in the above Corollary $E_1 = E = K$, $e_1 = \ldots = e_s = m_1 = \ldots = m_s = l_1 = \ldots = l_n = 0$, $n = 1$, $x_i = t_i$, $x = z$, $y_1 = \theta$, $z_1 = y \in \mathbb{Z}_p$,

$\delta \in \mathbb{Z}$, we get that the following sum converges in $\mathbb{T}_P$ and is in fact an entire function on $\mathbb{C}_P^{s+1}$:

$$\sum_{m\geq 0}\Big(\sum_{a\in A_+,\deg_\theta a=m,a\not\equiv 0 \pmod P}\omega_P(a)^\delta\frac{a(t_1)\cdots a(t_s)}{\langle a\rangle_P^y}\Big)z^m.$$

In particular, $\forall n\in\mathbb{Z}$, the sum

$$L_P(n;\underline{t};z):=\sum_{m\geq 0}\Big(\sum_{a\in A_+,\deg_\theta a=m,a\not\equiv 0 \pmod P}\frac{a(t_1)\cdots a(t_s)}{a^n}\Big)z^m\in\mathbb{T}_P$$

defines an entire function on $\mathbb{C}_P^{s+1}$. We refer the reader to [7], [10], for various arithmetic properties of "special values" of the series $L_P(n;\underline{t};z), n\in\mathbb{Z}$.

5.2. **Twisted Goss zeta functions.**

In this example, A is "general". Recall that $\mathbb{C}_\infty$ be the completion of $\overline{K}_\infty$. Let $s\geq 1$ be an integer. Let $\mathbb{F}\subset\overline{\mathbb{F}}_q$ be a field containing $\mathbb{F}_\infty$. Let $k_s(\mathbb{F})$ be defined as follows:
- if $s=1$, $k(\mathbb{F})=\mathbb{F}((y_1))$ where y_1 is an indeterminate over $\mathbb{F}$,
- if $s\geq 2$, let y_s be an indeterminate over $k_{s-1}(\mathbb{F})$, and set $k_s(\mathbb{F})=k_{s-1}(\mathbb{F})((y_s))$.
Observe that $\mathbb{F}$ is algebraically closed in $k_s(\mathbb{F})$.

Let L/K_∞ be a finite extension and let O_L be the valuation ring of L. Then:

$$O_L=\mathbb{F}_L[[\pi_L]],$$

where $\mathbb{F}_L$ is the residue field of L, and π_L is a prime of O_L. Let's consider the following tensor product:

$$k_s(\mathbb{F}_\infty)\otimes_{\mathbb{F}_\infty}O_L.$$

This ring can be identified naturally with $k_s(\mathbb{F}_L)\otimes_{\mathbb{F}_L}O_L$. Any element $f\in k_s(\mathbb{F}_L)\otimes_{\mathbb{F}_L}O_L$ can be written in a unique way:

$$\sum_{i\geq 0}\alpha_i\otimes\pi_L^i,\quad \alpha_i\in k_s(\mathbb{F}_L).$$

We set:

$$v_\infty(f)=\mathrm{Inf}\{v_\infty(\pi_L)i, i\in\mathbb{N},\alpha_i\neq 0\}.$$

Then v_∞ is a valuation on $k_s(\mathbb{F}_\infty)\otimes_{\mathbb{F}_\infty}O_L$ which does not depend on the choice of π_L. We denote by $k_s(\mathbb{F}_\infty)\widehat{\otimes}_{\mathbb{F}_\infty}O_L$ the completion of $k_s(\mathbb{F}_\infty)\otimes_{\mathbb{F}_\infty}O_L$ for v_∞. Finally denote by L_s the completion (for v_∞) of the field of fractions of $k_s(\mathbb{F}_\infty)\widehat{\otimes}_{\mathbb{F}_\infty}O_L$. If we identify $1\otimes\mathbb{F}_L$ with $\mathbb{F}_L$, and $1\otimes\pi_L$ with π_L which is thus an indeterminate over $k_s(\mathbb{F}_L)$, we have:

$$L_s=k_s(\mathbb{F}_L)((\pi_L)).$$

If $L\subset L'$, then we have a natural injective map compatible with v_∞:

$$L_s\hookrightarrow L'_s.$$

We denote by $\mathbb{C}_{\infty,s}$ the completion (for v_∞) of the inductive limit $\varinjlim_{L/K_\infty\text{ finite}}L_s$. Note that $\mathbb{C}_\infty, k_s(\overline{\mathbb{F}_q})\subset\mathbb{C}_{\infty,s}$, and the residue field of $\mathbb{C}_{\infty,s}$ is $k_s(\overline{\mathbb{F}_q})$.

Let z be an indeterminate over $\mathbb{C}_{\infty,s}$, we denote by $\mathbb{T}_z(\mathbb{C}_{\infty,s})$ the Tate algebra in the variable z with coefficients in $\mathbb{C}_{\infty,s}$.

We fix a K-embedding of $\overline{K}$ in $\mathbb{C}_\infty$. As in section 2, we consider [.] the Goss admissible map, i.e. for $I \in \mathcal{I}(A)$, we have:

$$[I] = \langle I \rangle \pi^{\frac{-\deg I}{d_\infty}} \in \pi^{\mathbb{Q}} U_\infty,$$

where π is our fixed prime of K_∞. Note that $V = K([I], I \in \mathcal{I}(A))$ is a finite extension of K. Recall that V is viewed as a subfield of $K_\infty([I], I \in \mathcal{I}(A))$. Let O_V be the integral closure of A in V. Since, for $a \in K^\times$, we have $[aA] = a/\operatorname{sgn}(a)$, we deduce that if I is a non-zero ideal of A then $[I] \in O_V$. Also recall that $\mathbb{F}_\infty \subset O_V$ is algebraically closed in V.

Let $s \in \mathbb{N}$. For $i = 1, \dots, s$, let $\rho_i : K_\infty([I], I \in \mathcal{I}(A)) \to \mathbb{F}_\infty((y_i))$ be a continuous $\mathbb{F}_p$-algebra homomorphism. Let $\mathbb{T}_{\underline{\rho},z}(\mathbb{C}_\infty)$ be the closure of $\mathbb{C}_\infty[z][\rho_i(O_V), i = 1, \dots, s]$ in $\mathbb{T}_z(\mathbb{C}_{\infty,s})$. Let also $\mathbb{T}_{\underline{\rho}}(\mathbb{C}_\infty)$ be the closure of $\mathbb{C}_\infty[\rho_i(O_V), i = 1, \dots, s]$ in $\mathbb{C}_{\infty,s}$.

Let's observe that $\mathbb{T}_{\underline{\rho},z}(\mathbb{C}_\infty)$ is an affinoid algebra over $\mathbb{C}_\infty$ (this is of also the case for $\mathbb{T}_{\underline{\rho}}(\mathbb{C}_\infty)$). Indeed, select $\theta \in A \setminus \mathbb{F}_q$. Then there exist $v_1, \dots, v_r \in O_V$ such that:

$$O_V = \oplus_{j=1}^r \mathbb{F}_\infty[\theta] v_j.$$

For $i = 1, \dots, s$, let $t_i = \rho_i(\theta)$. Then $t_1, \dots, t_s, z$ are $s+1$ indeterminates over $\mathbb{C}_\infty$. Let $\mathbb{T}_{s,z}(\mathbb{C}_\infty)$ be the Tate algebra in the variables $t_1, \dots, t_s, z$ with coefficients in $\mathbb{C}_\infty$. Clearly:

$$\mathbb{T}_{s,z}(\mathbb{C}_\infty) \subset \mathbb{T}_z(\mathbb{C}_{\infty,s}),$$

$$\mathbb{T}_{\underline{\rho},z}(\mathbb{C}_\infty) = \sum_{1 \leq i_1, \dots, i_s \leq r} \mathbb{T}_{s,z}(\mathbb{C}_\infty) \rho_1(v_{i_1}) \cdots \rho_s(v_{i_s}).$$

If we apply Theorem 4.1, we get as a special case:

Corollary 5.4. *Let E/K be a finite extension. Let $x \in \mathbb{Z}_p$, the following sum converges in $\mathbb{T}_{\underline{\rho},z}(\mathbb{C}_\infty)$:*

$$\sum_{m \geq 0} \Big(\sum_{\substack{\mathfrak{I} \in \mathcal{I}(\mathcal{O}_E), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = m}} \rho_1([N_{E/K}(\mathfrak{I})]) \cdots \rho_s([N_{E/K}(\mathfrak{I})]) \langle N_{E/K}(\mathfrak{I}) \rangle^x z^m.$$

Furthermore, as a function in z, it defines an entire function on $\mathbb{C}_\infty$ with values in $\mathbb{T}_{\underline{\rho}}(\mathbb{C}_\infty)$.

5.3. A-harmonic series attached to some admissible maps.

In this example, we work in the case A general. Let $\eta : \mathcal{I}(A) \to \overline{K}_\infty^\times$ be an admissible map such that $n(\eta) \in \mathbb{Z}$. Recall that there exist an open subgroup of finite index $M(\eta) \subset K_\infty^\times$, and an element $\alpha(\eta) \in \overline{K}_\infty^\times$, such that:

$$\forall I \in \mathcal{I}(A), \forall a \in K^\times \cap M(\eta), \quad \eta(Ia) = \eta(I) \Big(\frac{a}{\operatorname{sgn}(a)}\Big)^{n(\eta)} \alpha(\eta)^{v_\infty(a)}.$$

Note that there exists an open subgroup of finite index $N \subset K_\infty^\times$, $N \subset M(\eta)$, such that:

$$\forall a \in K^\times \cap N, \quad \operatorname{sgn}(\alpha(\eta))^{v_\infty(a)} = 1.$$

Let $\chi : \mathcal{I}(A) \to \overline{K}_\infty^\times$ be the map defined by:

$$\forall I \in \mathcal{I}(A), \quad \chi(I) = \pi^{v_\infty(\eta(I)) + \frac{\deg I}{d_\infty}(n(\eta) + v_\infty(\alpha(\eta)))} \operatorname{sgn}(\eta(I)).$$

If we set:

$$\mathcal{P} = \{xA, x \in K^\times \cap N\}.$$

Then $\frac{\mathcal{I}(A)}{\mathcal{P}}$ is a finite abelian group and:

$$\forall I \in \mathcal{I}(A), \forall J \in \mathcal{P}, \quad \chi(IJ) = \chi(I).$$

Let $n \in \mathbb{Z}$. For $u = (x, y) \in \mathbb{C}_\infty^\times \times \mathbb{Z}_p =: \mathbb{S}_\infty$, let's set:

$$L_{\eta,A}(\chi^n; u) = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(A), I \subset A \\ \deg I = m}} \chi^n(I) \langle \eta(I) \rangle^{-y} x^{-m}.$$

Then, by Theorem 4.1, $L_{\eta,A}(\chi^n; .)$ converges on $\mathbb{S}_\infty$. Now, let $x \in \mathbb{C}_\infty^\times$, and observe that:

$$L_{\eta,A}(\chi^{-n}; (\pi^{\frac{-n}{d_\infty}(n(\eta)+v_\infty(\alpha(\eta))} x, n)) = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(A), I \subset A \\ \deg I = m}} \eta(I)^{-n} x^{-m}.$$

Thus, for all $n \in \mathbb{Z}$, the following function in the variable z is entire on $\mathbb{C}_\infty$:

$$Z_{\eta,A}(n; z) = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(A), I \subset A \\ \deg I = m}} \eta(I)^{-n} z^m.$$

Observe that, by Lemma 3.2, if $m\, n(\eta) \leq 0$, we have:

$$Z_{\eta,A}(m; z) \in K(\eta(I), I \in \mathcal{I}(A))[z].$$

Let's furthermore assume that η is an algebraic admissible map. Let $E = K(\eta(I), I \in \mathcal{I}(A))$ which is a finite extension of K, and let $E_\infty = E \otimes_K K_\infty$. Let z be an indeterminate over K_∞, following [10], we denote by $\mathbb{T}_z(E_\infty)$ the closure of $E_\infty[z]$ in $E \otimes_K \widetilde{K_\infty}$, where $\widetilde{K_\infty}$ denotes the completion of the field of fractions of the Tate algebra in the variable z with coefficients in K_∞. From the above discussion, we get:

Corollary 5.5. *Let $m \in \mathbb{Z}$. The following sum converges in $\mathbb{T}_z(E_\infty)$:*

$$\zeta_{\eta,A}(m; z) := \sum_{k \geq 0} \sum_{\substack{I \in \mathcal{I}(A), I \subset A \\ \deg I = m}} \eta(I)^{-m} z^k.$$

Furthermore, if $m\, n(\eta) \leq 0$, we have:

$$\zeta_{\eta,A}(m; z) \in E[z].$$

Let v be a finite place of K, and let P_v be the maximal ideal of A associated to v. Let $\mathcal{I}(P_v)$ be the group of fractional ideals of A which are relatively prime to P_v. Let $\mathbb{C}_v$ be the v-adic completion of $\overline{K}_v$. Let η be our admissible map introduced in the beginning of the paragraph, and we assume that η is algebraic. Let $\chi : \mathcal{I}(P_v) \to \overline{K}_v$ such that:

$$\forall I \in \mathcal{I}(P_v), \quad \chi(I) = \omega_v(\eta(I)) \pi_v^{v(\eta(I)) + \frac{\deg I}{d_\infty} v(\alpha(\eta))}.$$

Note that there exists an open subgroup of finite index $N' \subset K_\infty^\times$, $N' \subset M(\eta)$, such that:

$$\forall a \in K^\times \cap N', \quad \omega_v(\alpha(\eta))^{v_\infty(a)} = 1.$$

Then:

$$\forall I \in I(P_v), \forall J \in \mathcal{P}', \quad \chi(IJ) = \chi(I),$$

where

$$\mathcal{P}' = \{xA \in \mathcal{I}(P_v), x \equiv 1 \pmod{P_v}, x \in K^\times \cap N' \cap \operatorname{Ker} \operatorname{sgn}\}.$$

Let $n \in \mathbb{Z}$. For $u = (x, y) \in \mathbb{C}_v^\times \times \mathbb{Z}_p$, let's set:

$$L_{v,\eta,A}(\chi^n; u) = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(P_v), I \subset A \\ \deg I = m}} \chi^n(I) \langle \eta(I) \rangle_v^{-y} x^{-m}.$$

Then, by Theorem 4.4, $L_{v,\eta,A}(\chi^n; .)$ converges on $\mathbb{C}_v^\times \times \mathbb{Z}_p$. Now, let $x \in \mathbb{C}_v^\times$, and observe that:

$$L_{v,\eta,A}(\chi^{-n}; (\pi_v^{\frac{-n}{d_\infty} v(\alpha(\eta))} x, n)) = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(P_v), I \subset A \\ \deg I = m}} \eta(I)^{-n} x^{-m}.$$

Thus, for all $n \in \mathbb{Z}$, the following function in the variable z is entire on $\mathbb{C}_v$:

$$Z_{v,\eta,A}(n; z) = \sum_{m \geq 0} \sum_{\substack{I \in \mathcal{I}(P_v), I \subset A \\ \deg I = m}} \eta(I)^{-n} z^m.$$

By Lemma 3.2, if $m\, n(\eta) \leq 0$, then:

$$Z_{v,\eta,A}(m; z) \in K(\eta(I), I \in \mathcal{I}(A))[z].$$

In particular, if $P_v = \alpha A$, $\alpha \in M(\eta)$ (there exist infinitely many such maximal ideals by Chebotarev's density Theorem), then, for $m \in \mathbb{Z}$ we get:

$$Z_{v,\eta,A}(m; z) = \Big(1 - (\frac{\alpha}{\mathrm{sgn}(\alpha)})^{-m n(\eta)} \alpha(\eta)^{m \frac{\deg P_v}{d_\infty}} z^{\deg P_v}\Big) Z_{\eta,A}(m; z).$$

We refer the interested reader to a forthcoming work of the authors dedicated to the arithmetic of such A-harmonic series ([12]).

6. Multiple several variable twisted zeta functions

We briefly explain how the constructions in Section 4 can easily be generalized in the spirit of Thakur's construction of positive characteristic multiple zeta values; the reader interested by the arithmetic of multiple zeta values for $K = \mathbb{F}_q(\theta)$ is referred to (this list is not exhaustive): [4], [39],[40], [25], [26], [14], [15], [29]. We keep the notation of Section 4.

Let $r \in \mathbb{N}, r \geq 1$. Let $s_1, \ldots, s_r \in \mathbb{N}$ and $n_1, \ldots, n_r \in \mathbb{N} \setminus \{0\}$. Let (F, v_F) be a field which is complete with respect to a non-trivial valuation $v_F : F \to \mathbb{R} \cup \{+\infty\}$ and such that F is an $\overline{\mathbb{F}}_q$-algebra. We set:

$$\mathbb{S}_{F,(n_1,\ldots,n_r)} = (F^\times \times \mathbb{Z}_p^{n_1}) \times \cdots \times (F^\times \times \mathbb{Z}_p^{n_r}).$$

For $i = 1, \ldots, r$, let $\psi_{1,i}, \ldots, \psi_{s_i,i}$ be s_i admissible maps such that $n(\psi_{j,i}) \in \mathbb{N}$. For $i = 1, \ldots, r, j = 1, \ldots, s_i$, let $\rho_{j,i} : K(\psi_{j,i}) \to F$ be an $\mathbb{F}_p$-algebra homomorphism. For $i = 1, \ldots, r$, we set:

$$\forall I \in \mathcal{I}(A), \quad \underline{\rho}_i(\underline{\psi}_i(I)) = \prod_{j=1}^{s_i} \rho_{j,i}(\psi_{j,i}(I)) \in F.$$

Let E/K be a finite extension. Let $\mathcal{B} \subset O_E$ be a non-zero ideal. Let $v_1, \ldots, v_n$ be the places of E above ∞, and for $j = 1, \ldots, n$, let N_j be an open subgroup of finite index of $E_{v_j}^\times$ (E_{v_j} is the v_j-adic completion of E), and we set : $N = \prod_{j=1,\ldots,n} N_j$.

Let $\mathcal{S}$ be a finite set, possibly empty, of maximal ideal of O_E which are relatively prime to $\mathcal{B}$. For $i = 1, \dots, r$, let $\chi_i : \mathcal{I}(\mathcal{B}) \to F$ be a map such that:

$$\forall \mathfrak{I} \in \mathcal{I}_(\mathcal{B}), \forall \mathfrak{J} \in \mathcal{P}(\mathcal{B}, N), \quad \chi_i(\mathfrak{I}\mathfrak{J}) = \chi_i(\mathfrak{I}).$$

For $i = 1, \dots, r$, let $\eta_{1,i}, \dots, \eta_{n_i,i}$ be n_i admissible maps with values in $\overline{K}_\infty^\times$. For $i = 1, \dots, r, j = 1, \dots, n_i$, let $\sigma_{j,i} : K(\langle \eta_{j,i} \rangle) \to F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. For $i = 1, \dots, r$, let $u_i = (x_i; z_{1,i}, \dots, z_{n_i,i}) \in F^\times \times \mathbb{Z}_p^{n_i}$, for $I \in \mathcal{I}(A)$, we set:

$$I^{u_i}_{\underline{\sigma}_i, \underline{\eta}_i} = x_i^{\deg I} \prod_{j=1}^{n_i} \sigma_{j,i}(\langle \eta_{j,i}(I) \rangle)^{z_{j,i}} \in F^\times.$$

We fix $i \in \{1, \dots, r\}$. Let $d \geq 0$ be an integer, for $u_i \in F^\times \times \mathbb{Z}_p^{n_i}$, we set:

$$S_{d;\mathcal{S},O_E}(\underline{\rho}_i, \underline{\psi}_i; \underline{\sigma}_i, \underline{\eta}_i; \chi_i; u_i) = \sum_{\substack{\mathfrak{I} \in \mathcal{I}_{\mathcal{S}}(\mathcal{B}), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = d}} \chi_i(\mathfrak{I}) \underline{\rho}_i(\underline{\psi}_i(N_{E/K}(\mathfrak{I}))) N_{E/K}(\mathfrak{I})^{-u_i}_{\underline{\sigma}_i, \underline{\eta}_i}.$$

By the proof of Theorem 4.1 and Remark 4.2, we get:

Corollary 6.1. *Let $u = (u_1, \dots, u_r) \in \mathbb{S}_{F,(n_1,\dots,n_r)}$, then the following sums converge in F :*

$$\sum_{d \geq 0} S_{d;\mathcal{S},O_E}(\underline{\rho}_1, \underline{\psi}_1; \underline{\sigma}_1, \underline{\eta}_1; \chi_1; u_1) \sum_{d > d_2 > \dots > d_r \geq 0} \prod_{j=2}^{r} S_{d_j;\mathcal{S},O_E}(\underline{\rho}_j, \underline{\psi}_j; \underline{\sigma}_j, \underline{\eta}_j; \chi_j; u_j),$$

$$\sum_{d \geq 0} S_{d;\mathcal{S},O_E}(\underline{\rho}_1, \underline{\psi}_1; \underline{\sigma}_1, \underline{\eta}_1; \chi_1; u_1) \sum_{d \geq d_2 \geq \dots \geq d_r \geq 0} \prod_{j=2}^{r} S_{d_j;\mathcal{S},O_E}(\underline{\rho}_j, \underline{\psi}_j; \underline{\sigma}_j, \underline{\eta}_j; \chi_j; u_j).$$

We furthermore assume that $n(\eta_{j,i}) \in \mathbb{Z}, j = 1, \dots, n_i, i = 1, \dots, r$, and that all our admissible maps are algebraic. Let v be a finite place of K, and Let P_v be the maximal ideal of A corresponding to the finite place v.

Let's set:

$$\mathbb{S}_{v,F,(n_1,\dots,n_r)} = (F^\times \times \mathbb{Z}_p^{n_1} \times \mathbb{Z}^{n_1}) \times \dots \times (F^\times \times \mathbb{Z}_p^{n_r} \times \mathbb{Z}^{n_r}).$$

For $i = 1, \dots, s, j = 1, \dots, s_i$, let $\rho_{j,i} : K(\psi_i) \hookrightarrow F$ be an $\mathbb{F}_p$-algebra homomorphism. For $i = 1, \dots, r$, we set as above:

$$\forall I \in \mathcal{I}(A), \quad \underline{\rho}_i(\underline{\psi}_i(I)) = \prod_{j=1}^{s_i} \rho_{j,i}(\psi_{j,i}(I)) \in F.$$

For $i = 1, \dots, r$, $j = 1, \dots, n_i$, let $\sigma_{j,i} : K_v(\langle \eta_i \rangle_v) \hookrightarrow F$ be a continuous $\mathbb{F}_p$-algebra homomorphism. For $i \in \{1, \dots, r\}$, for $u_i = (x; z_1, \dots, z_{n_i}; \underline{\delta}_i) \in F^\times \times \mathbb{Z}_p^{n_i} \times \mathbb{Z}^{n_i}$, for $I \in \mathcal{I}(A)$, we set:

$$I^{u_i}_{\underline{\sigma}_i, \underline{\eta}_i} = x^{\deg I} \prod_{j=1}^{n_i} \omega_v(\eta_{j,i}(I))^{\delta_{j,i}} \prod_{j=1}^{n_i} \sigma_{j,i}(\langle \eta_{j,i}(I) \rangle_v^{z_j}) \in F^\times.$$

Let E/K be a finite extension, and let $\mathcal{B}, N, \chi_1, \dots, \chi_r$, and $\mathcal{S}$ as above. Let $\mathcal{S}_v$ be the union of S and the maximal ideals of O_E above P_v and that do not divide $\mathcal{B}$.

We fix $i \in \{1, \dots, r\}$. Let $d \geq 0$ be an integer, for $u_i \in F^\times \times \mathbb{Z}_p^{n_i} \times \mathbb{Z}^{n_i}$, we set:

$$S_{v;d;\mathcal{S},O_E}(\underline{\rho}_i, \underline{\psi}_i; \underline{\sigma}_i, \underline{\eta}_i; \chi_i; u_i) = \sum_{\substack{\mathfrak{I} \in \mathcal{I}_{\mathcal{S}}(\mathcal{B}), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = d}} \chi_i(\mathfrak{I}) \underline{\rho}_i(\underline{\psi}_i(N_{E/K}(\mathfrak{I}))) N_{E/K}(\mathfrak{I})^{-u_i}_{\underline{\sigma}_i, \underline{\eta}_i},$$

$$S_{v;d;\mathcal{S}_v,O_E}(\underline{\rho}_i, \underline{\psi}_i; \underline{\sigma}_i, \underline{\eta}_i; \chi_i; u_i) = \sum_{\substack{\mathfrak{I} \in \mathcal{I}_{\mathcal{S}_v}(\mathcal{B}), \mathfrak{I} \subset O_E \\ \deg N_{E/K}(\mathfrak{I}) = d}} \chi_i(\mathfrak{I}) \underline{\rho}_i(\underline{\psi}_i(N_{E/K}(\mathfrak{I}))) N_{E/K}(\mathfrak{I})^{-u_i}_{\underline{\sigma}_i, \underline{\eta}_i}.$$

By the proof of Theorem 4.4, we get:

Corollary 6.2. *Let $u = (u_1, \dots, u_r) \in \mathbb{S}_{v,F,(n_1,\dots,n_r)}$, then the following sums converge in F :*

$$\sum_{d \geq 0} S_{v;d;\mathcal{S}_v,O_E}(\underline{\rho}_1, \underline{\psi}_1; \underline{\sigma}_1, \underline{\eta}_1; \chi_1; u_1) \sum_{d > d_2 > \dots > d_r \geq 0} \prod_{j=2}^{r} S_{v;d_j;\mathcal{S},O_E}(\underline{\rho}_j, \underline{\psi}_j; \underline{\sigma}_j, \underline{\eta}_j; \chi_j; u_j),$$

$$\sum_{d \geq 0} S_{v;d;\mathcal{S}_v,O_E}(\underline{\rho}_1, \underline{\psi}_1; \underline{\sigma}_1, \underline{\eta}_1; \chi_1; u_1) \sum_{d \geq d_2 \geq \dots \geq d_r \geq 0} \prod_{j=2}^{r} S_{v;d_j;\mathcal{S},O_E}(\underline{\rho}_j, \underline{\psi}_j; \underline{\sigma}_j, \underline{\eta}_j; \chi_j; u_j).$$

Let's pursue our basic example 5.3. Let $\eta : \mathcal{I}(A) \to \overline{K}_\infty^\times$ be an admissible map such that $n(\eta) \in \mathbb{Z}$. Let $r \geq 1$ be an integer and let $z_1, \dots, z_r$ be r indeterminates over $\mathbb{C}_\infty$. Let $\underline{n} = (n_1, \dots, n_r) \in \mathbb{Z}^r$. Let's define $Z_{\eta,A}(\underline{n}; \underline{z}) \in K(\eta_i(I), I \in \mathcal{I}(A), i = 1, \dots, r)[z_2, \dots, z_r][[z_1]]$ to be the following sum:

$$\sum_{d \geq 0} \sum_{\substack{I_1, \dots, I_r \in \mathcal{I}(A) \\ I_1, \dots, I_r \subset A \\ d = \deg I_1 > \deg I_2 > \dots > \deg I_r \geq 0}} \frac{1}{\eta_1(I_1)^{n_1} \cdots \eta_r(I_r)^{n_r}} z_1^{\deg I_1} \cdots z_r^{\deg I_r};$$

let's also define $Z^*_{\eta,A}(\underline{n}; \underline{z}) \in K(\eta_i(I), I \in \mathcal{I}(A), i = 1, \dots, r)[z_2, \dots, z_r][[z_1]]$ as the following sum:

$$\sum_{d \geq 0} \sum_{\substack{I_1, \dots, I_r \in \mathcal{I}(A) \\ I_1, \dots, I_r \subset A \\ d = \deg I_1 \geq \deg I_2 \geq \dots \geq \deg I_r \geq 0}} \frac{1}{\eta_1(I_1)^{n_1} \cdots \eta_r(I_r)^{n_r}} z_1^{\deg I_1} \cdots z_r^{\deg I_r}.$$

Then, as in example 5.3, by Corollary 6.1, we deduce that $Z_{\eta,A}(\underline{n}; \underline{z})$ and $Z^*_{\eta,A}(\underline{n}; \underline{z})$ define entire functions on $\mathbb{C}_\infty^r$. Let's also observe that, by Lemma 3.2, if $n_1 n(\eta) \leq 0$, then:

$$Z_{\eta,A}(\underline{n}; \underline{z}), Z^*_{\eta,A}(\underline{n}; \underline{z}) \in K(\eta_i(I), I \in \mathcal{I}(A), i = 1, \dots, r)[z_1, z_2, \dots, z_r].$$

Let's observe that, if $\eta = [.]$, $A = \mathbb{F}_q[\theta]$, and $\pi = \frac{1}{\theta}$, we recover Thakur's multiple zeta values (as in example 5.1, one can also recover their deformations in Tate algebras treated in [29]):

$$Z_{\eta,A}(\underline{n}; \underline{z}) = \sum_{d \geq 0} \sum_{\substack{a_1, \dots, a_r \in A_+ \\ d = \deg a_1 > \dots > \deg a_r \geq 0}} \frac{1}{a_1^{n_1} \cdots a_r^{n_r}} \prod_{i=1}^{r} z_i^{\deg_\theta a_i} \in K[z_2, \dots, z_r][[z_1]].$$

Let's assume that η is algebraic. Let v be a finite place of K, and let P_v be the maximal ideal of A associated to v. Let $\mathcal{I}(P_v)$ be the group of fractional ideals of A which are relatively prime to P_v. Let $\mathbb{C}_v$ be the v-adic completion of $\overline{K}_v$. We define

$Z_{v,\eta,A}(\underline{n};\underline{z}) \in K(\eta_i(I), I \in \mathcal{I}(A), i = 1,\dots,r)[z_2,\dots,z_r][[z_1]]$ to be the following sum:

$$\sum_{d\geq 0} \sum_{\substack{I_1\in\mathcal{I}(P_v), I_2,\dots,I_r\in\mathcal{I}(A)\\ I_1,\dots,I_r\subset A\\ d=\deg I_1>\dots>\deg I_r\geq 0}} \frac{1}{\eta_1(I_1)^{n_1}\cdots\eta_r(I_r)^{n_r}} z_1^{\deg I_1}\cdots z_r^{\deg I_r};$$

we also define $Z^*_{v,\eta,A}(\underline{n};\underline{z}) \in K(\eta_i(I), I \in \mathcal{I}(A), i = 1,\dots,r)[z_2,\dots,z_r][[z_1]]$ to be:

$$\sum_{d\geq 0} \sum_{\substack{I_1\in\mathcal{I}(P_v), I_2,\dots,I_r\in\mathcal{I}(A)\\ I_1,\dots,I_r\subset A\\ d=\deg I_1\geq\dots\geq\deg I_r\geq 0}} \frac{1}{\eta_1(I_1)^{n_1}\cdots\eta_r(I_r)^{n_r}} z_1^{\deg I_1}\cdots z_r^{\deg I_r}.$$

Again, as in example 5.3, by Corollary 6.2, we deduce that $Z_{v,\eta,A}(\underline{n};\underline{z})$ and $Z^*_{v,\eta,A}(\underline{n};\underline{z})$ define entire functions on $\mathbb{C}_v^r$.

For example, if $\eta = [.]$, $A = \mathbb{F}_q[\theta]$, and $\pi = \frac{1}{\theta}$, we have:

$$Z_{v,\eta,A}(\underline{n};\underline{z}) = \sum_{d\geq 0} \sum_{\substack{a_1,\dots,a_r\in A_+, a_1\notin P_v\\ d=\deg a_1>\deg a_2>\dots>\deg a_r\geq 0}} \frac{1}{a_1^{n_1}\cdots a_r^{n_r}} \prod_{i=1}^{r} z_i^{\deg_\theta a_i}.$$

Note that, if $n_1,\dots,n_r \leq 0$, then:

$$Z_{v,\eta,A}(\underline{n};\underline{z}) \equiv Z_{\eta,A}(\underline{n};\underline{z}) \pmod{P^{n_1}A[z_1,\dots,z_r]}.$$

Furthermore, for $\underline{n}\in\mathbb{Z}^r$, we observe that $Z_{v,\eta,A}(\underline{n};\underline{z})$ is the P_v-adic limit of certain sequences $Z_{\eta,A}((m_k,n_2,\dots,n_r);\underline{z}) \in K[z_1,\dots,z_r]$, $m_k \leq 0$, where m_k is suitably chosen and m_k converges p-adically to n_1. Indeed, let $\mathbb{T}_{\underline{z}}(K_v)$ be the Tate algebra in the variables $z_1,\dots,z_r$, with coefficients in K_v equipped with the Gauss valuation associated to v and still denoted by v. Observe that, if $m \leq 0$, we have:

$$\deg_{z_1} Z_{\eta,A}((m,n_2,\dots,n_r);\underline{z}) \leq \frac{\ell_q(-m)}{q-1},$$

where $\ell_q(-m)$ is the sum of the q-adic digits of $-m$. Now, as in the proof of Theorem 4.4, for $k \geq 0$, let's select $-m_k \in \mathbb{N}$ such that:

i) $-m_k \equiv -n_1 \pmod{q^{k+1}\mathbb{Z}_p}$,
ii)$-m_k \equiv -n_1 \pmod{q^{\deg P_v}-1}$,
iii) $\ell_q(-m_k) \leq (k+\deg P_v)(q-1)$,
iv) $-m_k \geq q^{k+1}$.

For example, if $-n_1 = \sum_{i\geq 0} a_i q^i$, $a_i \in \{0,\dots,q-1\}$, select $\delta_k \in \{1,\dots,q^{\deg P_v}-1\}$ such that $-n_1 - \sum_{i=0}^{k} a_i q^i \equiv \delta_k \pmod{q^{\deg P_v}-1}$, and set:

$$-m_k = \sum_{i=0}^{k} a_i q^i + \delta_k q^{(k+1)\deg P_v}.$$

Then:

$$v(Z_{\eta,A}((m_k,n_2,\dots,n_r);\underline{z}) - \sum_{d=0}^{k+\deg P_v} \sum_{\substack{a_1,\dots,a_r\in A_+, a_1\notin P_v\\ d=\deg a_1>\dots>\deg a_r\geq 0}} \frac{1}{a_1^{n_1}\cdots a_r^{n_r}} \prod_{i=1}^{r} z_i^{\deg_\theta a_i})$$
$$\geq q^{k+1} - (\mid n_2 \mid + \dots + \mid n_r \mid)(\deg P_v + k).$$

Thus in $\mathbb{T}_{\underline{z}}(K_v)$:

$$Z_{v,\eta,A}(\underline{n};\underline{z}) = \lim_k Z_{\eta,A}((m_k,n_2,\dots,n_r);\underline{z}).$$

To our knowledge, and in the case $A = \mathbb{F}_q[\theta]$, these type of elements $Z_{v,\eta,A}(\underline{n};\underline{1}) \in K_v$ have not been studied so far, and it would be very interesting to obtain informations on such type of objects, especially in the spirit of [4].

References

[1] G. Anderson, Rank one elliptic A-modules and A-harmonic series, *Duke Mathematical Journal* **73** (1994), 491-542.

[2] G. Anderson, Log-Algebraicity of Twisted A-Harmonic Series and Special Values of L-series in Characteristic p, *Journal of Number Theory* **60** (1996), 165-209.

[3] G. Anderson, D. Thakur, Tensor powers of the Carlitz module and zeta values, *Annals of Mathematics* **132** (1990), 159-191.

[4] G. Anderson, D. Thakur, Multizeta values for $\mathbb{F}_q[t]$, their period interpretation, and relations between them, *International Mathematics Research Notices* no. 11 (2009), 2038-2055.

[5] B. Anglès, F. Pellarin, Functional identities for L-series values in positive characteristic, *Journal of Number Theory* **142** (2014), 223-251.

[6] B. Anglès, F. Pellarin, Universal Gauss-Thakur sums and L-series, *Inventiones mathematicae* **200** (2015), 653-669 .

[7] B. Anglès, L. Taelman, with an appendix by V. Bosser, Arithmetic of characteristic p special L-values, *Proceedings of the London Mathematical Society* **110** (2015), 1000-1032.

[8] B. Anglès, F. Pellarin, F. Tavares Ribeiro, with an appendix by F. Demeslay, Arithmetic of positive characteristic L-series values in Tate algebras , *Compositio Mathematica* **152** (2016),1-61.

[9] B. Anglès, F. Pellarin, F. Tavares Ribeiro, Anderson-Stark Units for $\mathbb{F}_q[\theta]$, arXiv: 1501.06804 (2015).

[10] B. Anglès, F. Tavares-Ribeiro, Arithmetic of function fields units, arXiv:1506.06286 (2015).

[11] B. Anglès, T. Ngo Dac, F. Tavares Ribeiro, Exceptional Zeros of L-series and Bernoulli-Carlitz Numbers (with an appendix by B. Anglès, D. Goss, F. Pellarin, F. Tavares Ribeiro), arXiv: 1511.06209 (2015) .

[12] B. Anglès, T. Ngo Dac, F. Tavares Ribeiro, Arithmetic of special values of twisted zeta functions, in preparation.

[13] G. Böckle, Global L-functions over functions fields, *Mathematische Annalen* **323** (2002), 737-795 .

[14] C.-Y. Chang, Linear independence of monomials of multizeta values in positive characteristic, *Compositio Mathematica* **150** (2014), 1789-1808.

[15] C.-Y. Chang, M. Papanikolas, J. Yu, An effective criterion for Eulerian multizeta values in positive characteristic, arXiv: 1411.0124 (2014).

[16] F. Demeslay, A class formula for L-series in positive characteristic, arXiv:1412.3704 (2014).

[17] J. Fang, Special L-values of abelian t-modules, *Journal of Number Theory* **147** (2015), 300-325.

[18] J. Fang, Equivariant Special L-values of abelian t-modules, arXiv: 1503.07243 (2015) .

[19] D. Goss, v-adic zeta functions, L-series and measures for function fields, *Inventiones mathematicae* **55** (1979), 107-116.

[20] D. Goss, On a new type of L function for algebraic curves over finite fields, *Pacific Journal of Mathematics* **105** (1983), 143-181.

[21] D. Goss, Basic Structures of Function Field Arithmetic, Springer, 1996.

[22] D. Goss, Zeta phenomenology, Noncommutative geometry, arithmetic, and related topics, Johns Hopkins University Press, Baltimore, 2011, 159-182.

[23] D. Goss, On the L-series of F. Pellarin, *Journal of Number Theory* **133** (2013), 955-962.

[24] V. Lafforgue, Valeurs spéciales de fonctions L en caractéristique p, *Journal of Number Theory* **129** (2009), 2600-2634.

[25] J. Lara Rodríguez, D. Thakur, Zeta-like multizeta values for $\mathbb{F}_q[t]$, *Indian Journal of Pure and Applied Mathematics* **45** (2014), 787-801.

[26] J. Lara Rodríguez, D. Thakur, Multizeta shuffle relations for function fields with non rational infinite place, *Finite Fields and their Applications* **37** (2016), 344-356.

[27] M. Papanikolas, Log-Algebraicity on Tensor Powers of the Carlitz Module and Special Values of Goss L-Functions, work in progress.

[28] F. Pellarin, Values of certain L-series in positive characteristic, *Annals of Mathematics* **176** (2012), 2055-2093 .
[29] F. Pellarin, A note on multiple zeta values in Tate algebras, arXiv:1601.07348 (2016).
[30] F. Pellarin, R. Perkins, On certain generating functions in positive characteristic, arXiv: 1409.6006 (2014).
[31] F. Pellarin, R. Perkins, On twisted A-harmonic sums and Carlitz finite zeta values, arXiv:1512.05953 (2015).
[32] R. Perkins, On Pellarin's L-series, *Proceedings of the American Mathematical Society* **142** (2014), 3355-3368.
[33] R. Perkins, An exact degree for multivariate special polynomials, *Journal of Number Theory* **142** (2014), 252-263.
[34] R. Perkins, Explicit formulae for L-values in positive characteristic, *Mathematische Zeitschrift* **278** (2014), 279-299.
[35] L. Taelman, A Dirichlet unit theorem for Drinfeld modules, *Mathematische Annalen* **348** (2010), 899-907.
[36] L. Taelman, Special L-values of Drinfeld modules, *Annals of Mathematics* **175** (2012), 369-391.
[37] L. Taelman, A Herbrand-Ribet theorem for function fields, *Inventiones mathematicae* **188** (2012), 253-275.
[38] D. Thakur, Function Field Arithmetic, World Scientific, 2004.
[39] D. Thakur, Relations between multizeta values for $\mathbb{F}_q[t]$, *International Mathematics Research Notices* no. 12 (2009), 2318-2346.
[40] D. Thakur, Shuffle relations for function field multizeta values, *International Mathematics Research Notices* no. 11 (2010), 1973-1980.

Université de Caen Normandie, Laboratoire de Mathématiques Nicolas Oresme, CNRS UMR 6139, Campus II, Boulevard Maréchal Juin, B.P. 5186, 14032 Caen Cedex, France.
E-mail address: bruno.angles@unicaen.fr,tuan.ngodac@unicaen.fr,floric.tavares-ribeiro@unicaen.fr